

66-604

3

SID 66-421

STATISTICAL EVALUATION OF
PROTON RADIATION FROM SOLAR FLARES
FINAL TECHNICAL REPORT
JPL CONTRACT 951293
29 JULY 1966



FACILITY FORM 802

N 67 13210	(THRU)
176	1
(PAGES)	(CODE)
OR 80531	29
(NASA CR OR TMX OR AD NUMBER)	(CATEGORY)

GPO PRICE \$ _____

CFSTI PRICE(S) \$ _____

Hard copy (HC) 5.00

Microfiche (MF) 1.00

653 July 65

NORTH AMERICAN AVIATION, INC.
SPACE and INFORMATION SYSTEMS DIVISION

This work was performed for the Jet Propulsion Laboratory,
California Institute of Technology, sponsored by the
National Aeronautics and Space Administration under
Contract NAS7-100.

SID 66-421

STATISTICAL EVALUATION OF
PROTON RADIATION FROM SOLAR FLARES

FINAL TECHNICAL REPORT

JPL CONTRACT 951293

29 JULY 1966



Prepared by

J. B. Weddell

J. B. Weddell
Principal Investigator
Space Sciences

J. W. Haffner

J. W. Haffner
Co-Investigator
Aerothermal and Power
Systems

Approved by

F. J. Morin

F. J. Morin
Acting Director
Space Sciences

NORTH AMERICAN AVIATION, INC.
SPACE and INFORMATION SYSTEMS DIVISION



FOREWORD

This report describes the technical methods and results of effort on Contract 951293 with Jet Propulsion Laboratory, California Institute of Technology.

The authors appreciate the benefit of discussions with E. P. Divita of Jet Propulsion Laboratory. The assistance of Dorothy Steelman with the literature survey, data processing, and preparation of illustrations is gratefully acknowledged. R. Howard of Mount Wilson Observatory kindly permitted the reproduction of solar magnetograms.

PRECEDING PAGE BLANK NOT FILMED.



SUMMARY

The occurrence probability of solar high energy proton radiation has been evaluated statistically as a function of radiation flux, solar activity level, and solar distance. The results are applicable to determining the solar proton radiation environment of spacecraft in interplanetary space between 1 and 2 astronomical units (AU) in the 1969 to 1975 time period.

Previous tabulations of solar proton event characteristics have suffered from incompleteness and inconsistencies. A literature survey permitted a new tabulation of solar flare times and positions, proton radiation event onset, rise and decay times, and proton fluxes above 10, 30, and 100 Mev at 1 AU. At least partial data of 76 proton radiation events were established from direct observations described in the literature. From these data, an analytical representation expressing the temporal variation of the proton flux as a function of proton energy and total flux per radiation event was derived. This representation was used to estimate flux data not available from experimental results. The annual fluxes and the frequency distribution of radiation events versus flux per event were then calculated.

To estimate the probability distribution of expected solar proton radiation levels from 1969 to 1975, the sunspot number in these years was predicted by extrapolating the Fourier transformation of the sunspot numbers from 1749 to 1964. The correlation coefficient of the annual proton flux versus energy with the sunspot numbers from 1956 to 1963 was calculated. It was then possible to calculate the most probable proton fluxes versus energy for each year from 1969 to 1975, as well as the confidence levels of these flux values. The occurrence frequency distribution of solar proton radiation events was also correlated with the sunspot number, and the results were used to predict the most probable number of proton events of various sizes for each year from 1969 to 1975.

The long-term average flux predictions were supplemented by establishing correlations between the occurrence probabilities of individual solar flares and observable predictable solar activity parameters. These parameters were (1) the area, duration, and luminosity of calcium plage regions; (2) the existence of stationary points of zero magnetic field in dipolar magnetic regions; and (3) decreases of the sea-level neutron flux that precede some solar flare proton radiation events.

Theoretical analyses were used to evaluate the temporal variation of the solar proton flux versus energy as a function of solar distance. These analyses assumed (1) acceleration of protons by hydromagnetic shock waves



induced in the solar corona by absorption of light emitted by a flare; (2) motion of protons in the quiescent interplanetary magnetic field; and (3) random walk of energetic protons captured in the disordered magnetic fields associated with local concentrations of plasma in interplanetary space.

Three aspects of the solar proton radiation environment permit evaluation of the probability distribution of the environment level experienced on a trajectory between 1 and 2 AU during the 1969 to 1975 time period. These aspects are the radiation event occurrence frequency distribution as a function of flux per event and stage of the sunspot number cycle, the analytical representation of the flux versus time and energy in an individual event at 1 AU, and the dependence of the environment on solar distance. Application of these aspects to mission environment analysis is illustrated for the example of an earth-Mars trajectory in 1971.

Recommendations for additional study include analysis of angular distributions of the radiation flux versus time and energy, evaluation of the environments of energetic solar electrons, alpha particles and heavy nuclei, and extension of the solar distance range to 0.3 and 5 AU.



CONTENTS

Section		Page
	SUMMARY	v
I	INTRODUCTION	1
	Previous Radiation Environment Models	1
	Program Objective	1
	Task Description	2
II	PROTON RADIATION EVENT TABULATION	3
	Literature Survey	3
	Data Tabulation	5
III	CORRELATIONS WITH SOLAR PARAMETERS	23
	Selection of Correlations	23
	Proton Flux—Sunspot Number	24
	Proton Event Number—Sunspot Number	34
	Flare-Active Solar Region	37
	Proton Event—Neutron Monitor Rate	43
IV	PROTON RADIATION OCCURRENCE PROBABILITY	45
	Model Event at 1 AU	45
	Event Frequency Distribution	58
	Dependence on Solar Distance	73
V	RADIATION ENVIRONMENT EVALUATION	93
	Application to Mars Mission	93
VI	CONCLUSIONS	99
	Recommendations	99
VII	REFERENCES	103
	APPENDIXES	
	A. BIBLIOGRAPHY	109
	B. FLARE-PLAGE CORRELATION TABLES	153
	C. COMPUTER PROGRAM	169



Figure		Page
28	Probability of Absorption of Proton in Plasma	82
29	Space-Time Distribution of Protons in Plasma	83
30	Distribution of Proton Escape Times from Plasma	84
31	Proton Flux Rate Versus Energy at 1 AU	85
32	Proton Flux Rate Versus Energy at 1.41 AU	86
33	Proton Flux Rate Versus Energy at 2 AU	87
34	Line of Force in Quiescent Interplanetary Magnetic Field	89
35	Minimum Proton Energy at Receiver Versus Flare Longitude	90
36	Proton Flux Versus Energy at 1 AU	91
C1	SPOTNO Listing	171

TABLES

Table		Page
1	Solar Flare and Proton Event Data	6
2	Summary of 12 November 1960 Solar Proton Event Data	20
3	Monthly Smoothed Wolf Sunspot Numbers at Maxima	29
4	Correlation of Proton Flux With Sunspot Number	32
5	Proton Event Frequency Versus Flux Above 30 Mev	35
6	Intervals of Parameters of Plage Regions	37
7	Flare Prediction Accuracy	40
8	Magnetic Field - Flare Association	41
9	Accuracy of Combined Flare Prediction Method	42
10	Monthly Probabilities of Event Size for 1956-1963	69
11	Distribution of Events by Month, 1942-1963	72
12	Trajectory Parameters for a 1971 Mission to Mars	93
13	Solar Proton Events Expected at 1 AU for 1969-1975	97



ILLUSTRATIONS

Figure		Page
1	Sunspot Number Frequency Spectrum	27
2	Observed and Predicted Sunspot Numbers	30
3	Distribution of Deviations of Predicted Maximum Sunspot Numbers	31
4	Annual Proton Fluxes Versus Sunspot Number	33
5	Proton Event Number and Sunspot Number, 1940-1964	36
6	Onset Plus Rise Times Weighted for Flux Above 30 Mev	47
7	Onset Plus Rise Times Weighted for Flux Above 100 Mev	48
8	Characteristic Decay Times Weighted for Flux Above 30 Mev	49
9	Characteristic Decay Times Weighted for Flux Above 100 Mev	50
10	Integral Fluxes of Several Proton Events	52
11	Peak Flux Rates of Several Proton Events	53
12	Integral Flux Versus Peak Flux Rate Above 10 Mev	54
13	Integral Flux Versus Peak Flux Rate Above 30 Mev	55
14	Integral Flux Versus Peak Flux Rate Above 100 Mev	56
15	Integral Flux Above 30 Mev Versus Onset Plus Rise Time	59
16	Integral Flux Above 30 Mev Versus Characteristic Decay Time	60
17	Integral Flux Above 100 Mev Versus Onset Plus Rise Time	61
18	Integral Flux Above 100 Mev Versus Characteristic Decay Time	62
19	Calculated Peak Flux Rate Versus Parameter A	63
20	Calculated Integral Flux Versus Parameter A	64
21	Calculated Onset Plus Rise Time Versus Parameter A	65
22	Calculated Characteristic Decay Time Versus Parameter A	66
23	Distributions of Integral Flux Per Event	67
24	Monthly Proton Event Occurrence Probability	70
25	Monthly Distribution of Events, Event Number and Integral Flux	71
26	Proton Source Distribution Due to Hydromagnetic Shock	76
27	Proton Trajectory in Undisturbed Interplanetary Magnetic Field	77



I. INTRODUCTION


PREVIOUS RADIATION ENVIRONMENT MODELS

The design and operational philosophy of space vehicles and their components must be based on knowledge or assumptions concerning the environments that they are expected to encounter. The environment of charged particles, especially protons, at high kinetic energies is one to which many subsystems are sensitive. The significant proton radiation beyond the geomagnetic field is that originating at the sun; only protons associated with solar activity are discussed in this report. It is generally impossible to define this environment exactly, and models must be developed which express the probability of encountering various proton radiation levels.

Early attempts to provide models of the proton radiation environment were directed at establishing an upper envelope of the proton flux versus energy associated with a single solar flare (Reference 1). Later, the time-dependent flux versus energy distributions of several flare-associated increases of the proton radiation level were used to construct models of the proton flux as a function of energy and time after the flare (Reference 2). These models were also nearly envelopes of actual solar proton radiation events and did not represent the frequency of occurrence of events of various sizes. The desirability of establishing total flux encounter probabilities led to models (Reference 3) that expressed these probabilities as functions of proton energy and mission duration. The dependence of the expected environment on the stage of general solar activity and on solar distance are of great importance in planning and implementing future planetary exploration missions, and have not previously received quantitative consideration.

PROGRAM OBJECTIVE

The objective of this program is to evaluate the environment of high-energy proton radiation associated with solar flares in interplanetary space at distances from 1 to 2 astronomical units (AU) from the sun in 1969 to 1975. The results will be applicable to spacecraft environmental analysis and design.



TASK DESCRIPTION

In order to accomplish the general program objective, this effort was organized into six tasks with the following objectives:

1. Perform a literature survey of solar flare events for which protons have been observed, and prepare a table of characteristics of these events.
2. Establish statistical correlations between solar flare proton events and predictable solar phenomena.
3. Determine the occurrence probability of solar flare proton radiation characteristics.
4. Evaluate the spatial dependence of solar flare high-energy proton radiation characteristics.
5. Establish a method of evaluating the solar flare proton environment in space, and illustrate the method by application to a spacecraft trajectory in interplanetary space.
6. Prepare and submit reports as required, and provide for reviews of progress.



II. PROTON RADIATION EVENT TABULATION

LITERATURE SURVEY

While galactic (cosmic) radiation has been studied since 1913, the earliest measurements that suggested that the sun was a source of energetic particle radiation were made in 1942. The issue remained in doubt until 1956, when the unusual solar proton event of February 23 convinced even the skeptics. Since that date, approximately 100 events in which high-energy (Mev) protons were emitted from the sun have been recorded.

In order to prepare a table of the dates and characteristics of these events, a literature search was undertaken. In the initial stage, this search was based upon previous tabulations of solar proton events (References 4, 5, 6, and 7). These references not only led to a number of additional references relevant to solar proton events, but they also provided the bulk of the information for the list of the characteristics of the observed proton events.

The second stage of the literature search involved examining the indexes of standard technical journals in the field. The journals examined were:

(United States and Canada)

Journal of Geophysical Research (formerly Journal of Terrestrial Magnetism)

The Physical Review

Physical Review Letters

Canadian Journal of Physics

The Astronomical Journal

The Astrophysical Journal

Journal of Atmospheric and Terrestrial Physics

Science

Journal of the Royal Astronomical Society of Canada



(Western Europe)

Planetary and Space Sciences

Nature

Il Nuovo Cimento

Comptes Rendus

Space Science Reviews

Astronautica Acta

Space Research (COSPAR)

Monthly Notices of the Royal Astronomical Society

(Eastern Europe-English translations only)

Soviet Physics-JETP

Proceedings of the USSR Academy of Sciences

Geomagnetism and Aeronomy

Soviet Physics - Doklady

(Abstract Journals)

Physics Abstracts

Scientific and Technical Aerospace Reports (STAR)

International Aerospace Abstracts

As expected, the Journal of Geophysical Research yielded the most relevant information, followed by The Physical Review. For the early events (prior to 1956), The Physical Review was almost the sole source of information. The abstract journals were invaluable for locating articles written in languages other than English and reports not published in one of the standard journals.

Each journal was searched, starting with the most recent (generally December 1965 or January 1966) issue and going backward until it was apparent that further search had a very low probability of yielding anything.



For many of the journals, this cutoff point was January 1956, since those that failed to report the event of 23 February 1956 would not be expected to contain articles on earlier solar proton events. The Astrophysical Journal, The Astronomical Journal, and The Physical Review were searched back to 1950. The 19 November 1949 event was responsible for the first articles, although subsequent articles reported events that had been observed as early as 1942.

The references were divided into two groups — those concerned primarily with one event were categorized by month and year of that event (See Appendix A); those that referred to several events were categorized by the characteristic emphasized. Categories selected were:

Prediction of solar flares

Origin of solar flares

Source of solar flare particles and solar radio bursts

Propagation of solar flare particles

Spatial distributions of solar flare particles

Cosmic ray effects

Energy spectra of solar flare particles

Polar cap absorptions

Geomagnetic disturbances related to solar flares

Ionization and reactions in the earth's atmosphere

Miscellaneous

These lists of references appear in Appendix A. To prevent duplications and to facilitate study of the material, each list of references is presented in chronological order according to publication date.

DATA TABULATION

The first sources of information used in the table of solar flare and proton radiation event data (Table 1) were previously prepared tables of a similar nature (References 3, 4, 6, 8, and 9). The initial list prepared from



Table 1. Solar Flare and Proton Event Data

Solar Flare Data						Solar Particle Data				
Date Event No.	Time	Lat.	Long.	Plage	Imp.	Minimum Energy (Mev)	t (hours)	τ (hours)	$\hat{\phi}$ (p/cm ² -sec)	$\int \phi$ (p/cm ²)
2-28-42 1 **	1200	N 07	E 04		3	10 30 100	12* 6-8*	25* 15*	5,000* 1,000* 200*	1 x 10 ⁹ * 1.5 x 10 ⁸ * 1.5 x 10 ⁷ *
3-7-42 2 **	0442	N 07	W 90		?	10 30 100	12* 8*	30* 16*	7,000* 1,500* 250*	1.5 x 10 ⁹ * 2 x 10 ⁸ * 2 x 10 ⁷ *
7-25-46 3 **	1620	N 29	E 15		3+	10 30 100	10-15* 8*	30* 17*	10,000* 2,000* 400*	2 x 10 ⁹ * 2.5 x 10 ⁸ * 3 x 10 ⁷ *
5-11-49 4 **	2010	S 20	E 12		3+	10 30 100	10* 5*	20* 12*	1,000* 200* 30*	2 x 10 ⁸ * 2 x 10 ⁷ * 2 x 10 ⁶ *
11-19-49 5 **	1034	S 05	W 74		3	10 30 100	12-15* 8-10*	30* 20*	12,000* 3,000* 600*	3 x 10 ⁹ 5 x 10 ⁸ * 5 x 10 ⁷ *
1-27-51 6 **						10 30 100	10-12* 6-8*	20-25* 12-15*	2,000* 300* 50*	3 x 10 ⁸ * 3 x 10 ⁷ * 3 x 10 ⁶ *
5-16-51 7 **	1702	N 15	W 05		2	10 30 100	10-12* 6-8*	25* 15*	3,000* 500* 100*	5 x 10 ⁸ * 5 x 10 ⁷ * 5 x 10 ⁶ *
1-17-52 8 **						10 30 100	10* 5-8*	20-25* 10-15*	1,500* 250* 40*	2 x 10 ⁸ * 3 x 10 ⁷ * 3 x 10 ⁶ *

*Estimated value (see Sections II and IV).

**Characteristics estimated from balloon-borne ionization chambers and/or sea level neutron monitor data.

***Probably no patrol.

*Estimated value (see Sections II and IV).

**Characteristics estimated from balloon-borne ionization chambers and/or sea level neutron monitor data.

***Probably no patrol.



Table 1. Solar Flare and Proton Event Data (Cont)

Solar Flare Data						Solar Particle Data				
Date Event No.	Time	Lat.	Long.	Plage	Imp.	Minimum Energy (Mev)	t (hours)	τ (hours)	$\hat{\phi}$ (p/cm ² -sec)	$\int \phi$ (p/cm ²)
2-23-56 9	0340	N 22	W 74	3400	3+	10 30 100	6-8 3-4	30 16	10,000 8,000 5,000	1.8×10^9 + 1.0×10^9 ✓ 3.5×10^8 □
8-31-56 10	1241	N 16	E 16	3643	3	10 30 100	15* 8-10*	36 20*	300* 150 60	8×10^7 * + 2.5×10^7 ✓ 6×10^6 □
11-13-56 11	1501	N 16	W 10	3753	2+	10 30 100	12* 7*	25* 16*	4,000* 800* 150*	8×10^8 * + 1.0×10^8 ✓ 1.5×10^7 * □
1-20-57 12	1120	S 25	W 30	3820	3+	10 30 100	10* 8*	24 18*	30,000* 2,000* 100	5×10^9 * 3×10^8 1×10^7
4-3-57 13	0835	S 15	W 60	3907	3	10 30 100	11* 7*	23* 15*	2,400* 450* 70*	4×10^8 * 5.0×10^7 5×10^6 *
6-21-57 14	1780*	N 14	E 02		2	10 30 100	12* 7*	26* 16*	5,000* 1,000* 200*	1.0×10^9 * 1.5×10^8 1.5×10^7 *
7-3-57 15	0740	N 14	W 40	4039	3+	10 30 100	2* 1-2*	5* 3*	20,000* 2,000 100	2×10^8 * 2×10^7 1×10^6 *



Table 1. Solar Flare and Proton Event Data (Cont)

Solar Flare Data						Solar Particle Data				
Date Event No.	Time	Lat.	Long.	Plage	Imp.	Minimum Energy (Mev)	t (hours)	τ (hours)	$\hat{\phi}$ (p/cm ² -sec)	$\int \phi$ (p/cm ²)
7-24-57 16	1827	S 24	W 27	4070	2	10 30 100	9* 5*	20* 12*	600* 100* 15*	8 x 10 ⁷ * 7.5 x 10 ⁶ 7.7 x 10 ⁵
8-9-57 17	0628	S 09	E 75	4099	2	10 30 100	8* 5*	18 12*	150* 20 2*	1.5 x 10 ⁷ * 1.5 x 10 ⁶ 1 x 10 ⁵ *
8-29-57 18	0555 1037	N 24 S 25	E 35 E 20	4124 4125	2 2	10 30 100	25* 15*	60 40*	10,000* 600 30	2 x 10 ⁹ * 1.2 x 10 ⁸ 3 x 10 ⁶
8-31-57 19	0552 0552 1312	S 31 N 13 N 25	W 02 E 03 W 02	4125 4124 4124	2+ 2 3	10 30 100	11* 7*	25* 15*	3,000* 600* 100*	6 x 10 ⁸ * 8 x 10 ⁷ 8 x 10 ⁶
9-2-57 20	1303 1316	N 10 S 34	W 26 W 36	4124 4125	1+ 2+	10 30 100	11* 7*	24* 15*	2,500* 400* 70*	4 x 10 ⁸ * 5.0 x 10 ⁷ 4.5 x 10 ⁶
9-12-57 21	1516	N 11	W 18	4134	2	10 30 100	8* 5*	18* 11*	400* 60* 8*	6 x 10 ⁷ * 6.0 x 10 ⁶ 5 x 10 ⁵
9-21-57 22	1338	N 10	W 06	4152	3	10 30 100	10* 7*	24 15-20*	600* 100 15*	1.5 x 10 ⁷ * 1.5 x 10 ⁶ 1 x 10 ⁵ *



Table 1. Solar Flare and Proton Event Data (Cont)

Solar Flare Data						Solar Particle Data				
Date Event No.	Time	Lat.	Long.	Plage	Imp.	Minimum Energy (Mev)	t (hours)	τ (hours)	$\hat{\phi}$ (p/cm ² -sec)	$\int \phi$ (p/cm ²)
9-26-57 23	2000*	N 22	E 15		3	10 30 100	7* 4*	15* 9*	50* 6* 1*	6 x 10 ⁶ * 4 x 10 ⁵ * 2.0 x 10 ⁴
10-20-57 24	1642	S 26	W 45	4189	2+	10 30 100	12* 8*	30 20*	1,500* 500 140	2 x 10 ⁸ * 5 x 10 ⁷ 1 x 10 ⁷
11-4-57 25	***					10 30 100	8* 5*	18 12*	350* 50 5*	1 x 10 ⁸ * 9 x 10 ⁶ 8 x 10 ⁵ *
2-9-58 26	2142	S 12	W 14	4404	2+	10 30 100	8* 5*	18 12*	5,000* 1,000 200*	5 x 10 ⁷ * 5 x 10 ⁶ 4 x 10 ⁵
3-23-58 27	1005	S 14	E 78	4476	3+	10 30 100	18* 8*	40 20	8,000 1,200 100	2 x 10 ⁹ 2.5 x 10 ⁸ 1 x 10 ⁷
3-25-58 28	0530 0603	N 17 S 15	E 25 E 50	4474 4476	2 2	10 30 100	14* 9*	30* 20*	18,000* 4,500* 900*	4 x 10 ⁹ * 6 x 10 ⁸ 7 x 10 ⁷ *
4-10-58 29	0100 0914	N 25 N 18	W 63 W 78	4485 4485	1 1+	10 30 100	8* 5*	18* 11*	350* 50 6*	5 x 10 ⁷ * 5 x 10 ⁶ 4 x 10 ⁵ *



Table 1. Solar Flare and Proton Event Data (Cont)

Solar Flare Data						Solar Particle Data				
Date Event No.	Time	Lat.	Long.	Plage	Imp.	Minimum Energy (Mev)	t (hours)	τ (hours)	$\hat{\phi}$ (p/cm ² -sec)	$\int \phi$ (p/cm ²)
7-7-58 30	0115	N 25	W 08	4634	3+	10 30 100	15* 6-10*	32 16	10,000 1,500 100	1.8 x 10 ⁹ 2.5 x 10 ⁸ 9 x 10 ⁶
7-29-58 31	0056 0303	S 17 S 14	W 42 W 44	4659 4659	2+ 3	10 30 100	9* 5-6*	20* 12*	550* 90* 10*	8 x 10 ⁷ * 8.5 x 10 ⁶ 7 x 10 ⁵
8-16-58 32	0440	S 14	W 50	4686	3+	10 30 100	10 6*	18 14*	3,000 400 20	4 x 10 ⁸ 4 x 10 ⁷ 1.6 x 10 ⁶
8-21-58 33	1927	N 08	E 59		1	10 30 100	6-7* 4*	14* 9*	40* 5* <1*	4 x 10 ⁶ * 3 x 10 ⁵ * 1.9 x 10 ⁴
8-22-58 34	1448	N 18	W 10	4708	3	10 30 100	10 3	20 8-12	6,000 600 20	8 x 10 ⁸ 7 x 10 ⁷ 1.8 x 10 ⁶
8-26-58 35	0027	N 20	W 54	4708	3	10 30 100	9 5*	12 7*	15,000 1,500 50	1.5 x 10 ⁹ 1.1 x 10 ⁸ 2.0 x 10 ⁶
9-22-58 36	0750 1017	S 19 N 17	W 42 W 65	4765 4756	2 2	10 30 100	15* 8*	30 16*	600 50 1.5	9 x 10 ⁷ 6 x 10 ⁶ 1 x 10 ⁵



Table 1. Solar Flare and Proton Event Data (Cont)

Solar Flare Data						Solar Particle Data				
Date Event No.	Time	Lat.	Long.	Plage	Imp.	Minimum Energy (Mev)	t (hours)	τ (hours)	$\hat{\phi}$ (p/cm ² -sec)	$\int \phi$ (p/cm ²)
5-10-59 37	2140	N 18	E 48	5148	3+	10 30 100	20 12	22 14	30,000 6,000 1,000	5.5×10^9 9.6×10^8 8.5×10^7
6-13-59 38	0400	N 17	E 57	5204	1+	10 30 100	20* 12*	40* 24	80* 9 1*	1×10^9 * 8.5×10^7 7×10^6 *
7-10-59 39	0240	N 20	E 66	5270	3+	10 30 100	30 18	40 20	15,000 4,000 1,200	4.5×10^9 1.0×10^9 1.0×10^8
7-14-59 40	0349	N 16	E 07	5269	3+	10 30 100	16 12	18 12	50,000 10,000 1,200	7.5×10^9 1.3×10^9 1.0×10^8
7-16-59 41	2145 2132	N 08 N 15	W 26 W 30	5269	3+ 3	10 30 100	12 4	30 18	18,000 6,000 1,500	3.3×10^9 9.1×10^8 1.3×10^8
8-18-59 42	1030	N 12	W 33		3	10 30 100	27* 24	30* 25*	150* 20 2*	2×10^7 * 1.8×10^6 1×10^5 *
9-2-59 43	2000*	N 12	E 60			10 30 100	9* 6*	20* 13*	800* 150* 20*	1×10^8 * 1.2×10^7 1×10^6 *



Table 1. Solar Flare and Proton Event Data (Cont)

Solar Flare Data						Solar Particle Data				
Date Event No.	Time	Lat.	Long.	Plage	Imp.	Minimum Energy (Mev)	t (hours)	τ (hours)	$\hat{\phi}$ (p/cm ² -sec)	$\int \phi$ (p/cm ²)
1-11-60 44	2100	N 23	E 03		3	10 30 100	6 4*	30 20*	25* 2.5 0.2*	6×10^6 * 4×10^5 2.5×10^4 *
3-29-60 45	0800*	N 11	E 30			10 30 100	8* 5*	18* 11*	400* 60* 8*	6×10^7 * 6.0×10^6 5×10^5 *
3-30-60 46	1540	N 12	E 12	5615	2	10 30 100	7* 4*	18* 10*	500* 60* 6*	7×10^7 * 6.0×10^6 4×10^5
4-1-60 47	0858	N 12	W 10	5615	3	10 30 100	2 1	6-12 4	400 200 45	1.5×10^7 5.0×10^6 8.5×10^5
4-5-60 48	0245	N 10	W 61	5615	2+	10 30 100	3 2*	12 8*	150* 20 2*	1.4×10^7 1.1×10^6 7×10^4 *
4-28-60 49	0130	S 05	E 34	5645	3	10 30 100	3 2	8 5	250 120 25	1.3×10^7 5.0×10^6 7×10^5
4-29-60 50	0220 0420 0554	N 12 N 10 N 14	W 20 W 22 W 20	5642 5642 5642	2 3 2	10 30 100	9* 5*	20* 11*	600* 100 15*	7×10^7 * 7×10^6 6×10^5 *



Table 1. Solar Flare and Proton Event Data (Cont)

Solar Flare Data						Solar Particle Data				
Date Event No.	Time	Lat.	Long.	Plage	Imp.	Minimum Energy (Mev)	t (hours)	τ (hours)	$\hat{\phi}$ (p/cm ² -sec)	$\int \phi$ (p/cm ²)
5-4-60 51	1020	N 14	W 90	5642	3+	10 30 100	2-3 1	8-14 4-6	150 100 40	1.2×10^7 6×10^6 1.2×10^6
5-6-60 52	1450	S 10	E 08	5653	3	10 30 100	3 2*	24 15*	300* 40 4*	4×10^7_6 4×10^6 $3 \times 10^5_*$
5-13-60 53	0536	N 30	W 64	5654	3+	10 30 100	3 2*	15 10*	180 60 10	1.5×10^7 4×10^6_5 4.5×10^5
6-1-60 54	0844	N 28	E 46		3+	10 30 100	2 1.5*	20 12*	50* 5 0.4*	$5 \times 10^6_*$ 4×10^5 $3 \times 10^4_*$
8-12-60 55	1940	N 22	E 27		3+	10 30 100	10 7*	50 35*	25* 3 0.3*	$8 \times 10^6_*$ $6 \times 10^5_*$ $4 \times 10^4_*$
9-3-60 56	0110	N 18	E 88	5838	3	10 30 100	12-16 6-9	32 26	450 200 60	9×10^7 3.5×10^7 7×10^6
9-26-60 57	0550	S 21	W 64	5858	2	10 30 100	5* 3*	10* 7*	300 40 4	2×10^7 2×10^6 1.2×10^5



Table 1. Solar Flare and Proton Event Data (Cont)

Solar Flare Data						Solar Particle Data				
Date Event No.	Time	Lat.	Long.	Plage	Imp.	Minimum Energy (Mev)	t (hours)	τ (hours)	$\hat{\phi}(\text{p}/\text{cm}^2\text{-sec})$	$\int \phi(\text{p}/\text{cm}^2)$
11-12-60 58	1329	N 27	W 02	5925	3+	10 30 100	10-16 8-10	18-24 14-18	32,000 12,000 2,500	4×10^9 1.3×10^9 2.5×10^8
11-15-60 59	0221	N 30	W 32	5925	3+	10 30 100	8-16 3-5	16-20 8-12	22,000 8,000 2,400	2.5×10^9 7.2×10^8 1.2×10^8
11-20-60 60	2020	N 28	W 113	5925	3	10 30 100	2-4 1	10-16 4-6	2,200 1,000 400	1.4×10^8 4.5×10^7 8×10^6
7-11-61 61	1700	S 06	E 32	6171	3	10 30 100	8-10 4	22-26 18	120 30 3	1.7×10^7 3×10^6 3×10^5
7-12-61 62	1030	S 07	E 22	6171	3+	10 30 100	8-12 6	16-20 12	4,000 400 15	5×10^8 4×10^7 1×10^6
7-15-61 63	1440 1600 1512	N 14 S 07	E 13 W 20	6172 6172	3 2	10 30 100	9* 5*	20* 12*	800* 150* 20*	$1 \times 10^{8*}$ 1.3×10^7 1×10^6
7-18-61 64	1010	S 06	W 60	6171	3+	10 30 100	6-10 2-3	24 12	7,000 2,500 600	1×10^9 3×10^8 4×10^7



Table 1. Solar Flare and Proton Event Data (Cont)

Solar Flare Data						Solar Particle Data				
Date Event No.	Time	Lat.	Long.	Plage	Imp.	Minimum Energy (Mev)	t (hours)	τ (hours)	ϕ (p/cm ² -sec)	$\int \phi$ (p/cm ²)
7-20-61 65	1620 1830	S 07	W 90	6171	3+	10 30 100	4-6 1.5	6-8 4	400 150 40	1.5×10^7 5×10^6 9×10^5
7-28-61 66	1920* 2180* 2145*	S 18 N 53 N 16	W 90 W 90 W 90	6175	1 1 1	10 30 100	8* 5*	18* 11*	400* 60* 7*	5×10^7 * ₆ 4.4×10^6 4×10^5 *
9-8-61 67	2200*	N 24	W 90		1	10 30 100	8* 5*	17* 11*	250* 40* 4*	3×10^7 * 3.0×10^6 3×10^5 *
9-10-61 68	2020	N 12	W 90	6212	1	10 30 100	10 6-7*	23* 14*	2,000* 350* 60*	3×10^8 * 4×10^7 4×10^6 *
9-28-61 69	2223	N 14	E 30	6235	3	10 30 100	1.5 1	12 8	500 150 36	5×10^7 6×10^6 1.1×10^6
11-10-61 70	1444	N 08	W 90	6264	2+	10 30 100	9* 5-6*	19* 12*	600* 90* 10*	8×10^7 * 8×10^6 7×10^5 *



Table 1. Solar Flare and Proton Event Data (Cont)

Solar Flare Data						Solar Particle Data				
Date Event No.	Time	Lat.	Long.	Plage	Imp.	Minimum Energy (Mev)	t (hours)	τ (hours)	$\hat{\phi}$ (p/cm ² -sec)	$\int \phi$ (p/cm ²)
2-23-62 71	1848	S 09	E 29	6351	1+	10 30 100	2* 1*	4* 2.5*	15* 1* <1*	1 x 10 ⁶ * 8 x 10 ⁴ * 4 x 10 ³ *
10-23-62 72	1708	N 02	W 71		2	10 30 100	2 1.5*	8 6*	16 4.5 0.5	6 x 10 ⁵ 1.2 x 10 ⁵ 1 x 10 ⁴
2-9-63 73	***					10 30 100	3* 2*	7* 5*	6* 1* <1*	3 x 10 ⁵ * 2 x 10 ⁴ * 1 x 10 ³ *
4-15-63 74	1620	S 09	W 09	6766	2+	10 30 100	7* 4*	16* 10*	25* 2* <1*	5 x 10 ⁶ * 4 x 10 ⁵ * 3 x 10 ⁴ *
9-20-63 75	2351	N 10	W 09	6964	2+	10 30 100	12* 8*	30* 18*	10,000* 2,500* 500*	2 x 10 ⁹ * 3 x 10 ⁸ * 4 x 10 ⁷ *
9-26-63 76	0701	N 13	W 76	6964	3	10 30 100	10* 7*	25* 15*	3,000* 500* 100*	5 x 10 ⁸ * 6 x 10 ⁷ * 6 x 10 ⁶ *



these sources included some or all of the following data for each event and the flare most probably associated with each event:

Flare Data

Date of occurrence

Universal time of maximum

Heliographic latitude and longitude

Calcium plage region number, assigned by the McMath-Hulbert Observatory

Importance (1 - to 3 +)

Proton Radiation Event Data*

Onset plus rise time (t) in hours (from optical flash maximum to peak proton flux)

Characteristic decay time (τ) in hours (time in which proton flux decreases from peak to $1/e$ of peak)

Peak flux rate ($\hat{\phi}$) in $\text{p/cm}^2 \text{ sec}$ above energy E_0 (Mev)

Integral flux ($\int \phi$) in p/cm^2 above energy E_0 (Mev)

The solar proton data are generally given for particles above energies of 10, 30 and 100 Mev. However, frequently a complete set of the data for a given event (especially the early events) was not available. In particular, values of t (the onset plus rise time) and τ (the characteristic decay time) for protons above 10 Mev were not available for any of the events.

A secondary source of proton event information was Reference 10, which lists the integral fluxes for protons above 30 Mev for 52 events from 23 February 1956 to 10 November 1961. Of these events, fluxes for the following dates were not reported in the other compilations examined: 13 November 1956, 3 April 1957, 21 June 1957, 25 March 1958, 2 September 1959, 29 March 1960, 15 July 1961, 28 July 1961, 8 September 1961, 10 September 1961 and 10 November 1961.

Another very useful source of information for the early events was Reference 11, which, in addition to listing and providing information on nine events from 1947 to 1952, referenced earlier work pertaining to solar proton

*As observed at earth (1 AU)



events. Based on these articles, the following events were selected because the measurements (either by balloon-borne detectors or ground-based neutron monitors) indicated the presence of solar protons: 28 February 1942, 7 March 1942, 25 July 1946, 11 May 1949, 19 November 1949, 27 January 1951, 16 May 1951 and 17 January 1952. A problem arose in that the information desired for the table was not available directly. However, by comparing neutron monitor data (principally from Deep River, Canada) and ionization detector measurements (principally at Cheltenham, Maryland; Climax, Colorado; and Godhavn, Greenland) for these events with those for later events, estimates of the integral proton fluxes above 100 Mev were made.

The late 1962 and 1963 events presented a different kind of problem. The time required for data to be analyzed and reported in the literature appears to run from one to two years. Therefore, the only article of significance located for these events was that written by Geodeke and Masley (Reference 12). They report riometer attenuations at 30 Mcs for the following events: 23 February 1962, 23 October 1962, 9 February 1963 and 15 April 1963. The peak flux rate above 10 Mev was related to the maximum riometer attenuation by

$$\hat{\phi} > 10 \text{ Mev (p/cm}^2\text{-sec)} = 30 a^2 \quad (1)$$

where a is the attenuation in db. This relationship, obtained by fitting the data from Webber's table (Reference 4), is a reasonable average, although a factor of 3 variation is observed. By using this relationship, the peak flux rates above 10 Mev for these four events were estimated.

After these data were gathered, two problems presented themselves. One problem was to estimate the incomplete data for several of the events; the other problem was to reconcile differences in the data for those events which were reported by more than one source. The estimation of the missing data was carried out by use of the model event described in Section IV. Basically, this involved obtaining mathematical relationships that are reasonable approximations for those events for which all or most of the data are available and using these relationships to estimate the missing data from the available data. In this way, the entire event could be approximately reconstructed once a single flux, flux rate, rise, or decay time was known (provided the energy range represented was known). All of the missing data were estimated in this way. The estimated numbers are followed by an asterisk (*) in Table 1. The reconciliation of conflicts was sometimes carried out by use of the model event also. If a decision could not be made on the basis of reliability of measurement, reputation of author, or date of publication (later publications were favored), the data that best fit the other data (on the basis of the model) were used. The numbers finally selected are listed in Table 1.



In order to illustrate the techniques that were used to arrive at the numbers in Table 1, the event of 12 November 1960 will be considered. This event has been reported by many authors and discussed in some detail by Webber (Reference 13), Fitchel, Guss, and Ogilvie (Reference 14), Lewis et al (Reference 6), and several others. Unfortunately, the available measurements have not been collected by holding a symposium on the event, as was done for the July 1959 events (Reference 15).

One problem is to distinguish between primary references (in which authors report the results of direct measurements) and secondary references (in which authors report event characteristics based upon information obtained from the literature, private communication with other workers in the field, etc.). For example, various authors have used Webber's tabulations in their own publications. While secondary references are quite valuable in a study of this sort, it is important to prevent an artificial weighting of the results by citing a number of secondary references based upon one or a few primary references.

A brief summary of the reported characteristics of the event of 12 November 1960 is given in Table 2. These are either primary references or authoritative secondary references based upon a study of the available primary references. An integral flux of $1.2 \times 10^9 \text{ p/cm}^2 > 30 \text{ Mev}$ appears to be a common meeting ground, with the values of $\sim 2 \times 10^8$ being in units of $\text{p/cm}^2\text{-ster}$. Similarly, an integral flux of $\sim 3.5 \times 10^8 \text{ p/cm}^2 > 100 \text{ Mev}$ agrees with the published numbers. In a similar manner, peak flux rates of $12,000 \text{ p/cm}^2\text{-sec} (> 30 \text{ Mev})$ and $2,500 \text{ p/cm}^2\text{-sec} (> 100 \text{ Mev})$ were arrived at. Rise times of 10 to 16 hr ($> 30 \text{ Mev}$) and 8 to 10 hr ($> 30 \text{ Mev}$) and characteristic decay time of 18 to 24 hr ($> 30 \text{ Mev}$) and 14 to 18 hr ($> 100 \text{ Mev}$) were selected in a similar way.

In addition to the numbers reported in Table 1, several other references reporting characteristics of the 12 November 1960 event were found. Spectral shape information has been reported by de Feiter, Freon, and Le Grand (Reference 25), Kodama and Kitamura (Reference 26), Pomerantz, Duggal, and Nagashima (Reference 27), Conforto and Iucci (Reference 28), Lockwood and Shea (Reference 29), Roederer et al (Reference 30), and Ogilvie and Bryant (Reference 31). These references show that there seems to be a common meeting ground in the vicinity of an R^{-6} spectrum, although several authors point out that the spectral shape appeared to be too complex to be fit by a simple power law, either in rigidity or energy. The double peak in the neutron monitor data, reaching 2.1 times the normal sea-level counting rate, probably was related to this fact. The second peak has been interpreted as due to the influence of disturbed magnetic fields which "channeled" the solar plasma cloud to earth.



Table 2. Summary of 12 November 1960 Solar Proton Event Data

Ref.	t (hr)	τ (hr)	$\hat{\phi}$ (p/cm ² -sec)	$\int \phi p / \text{cm}^2$
9	12-16 (>30 Mev) 8-10 (>100 Mev)	18-24 (>30 Mev) 14-18 (>100 Mev)	12,000 (>30 Mev) 2,500 (>100 Mev)	1.4 x 10 ⁹ (>30 Mev) 3.5 x 10 ⁸ (>100 Mev)
4	10 (>30 Mev) 8 (>100 Mev)	18 (>30 Mev) 14 (>100 Mev)	12,000 (>30 Mev) 2,500 (>100 Mev)	1.3 x 10 ⁹ (>30 Mev) 2.5 x 10 ⁸ (>100 Mev)
16				3.9 x 10 ⁸ (>100 Mev) (includes 12-15 Nov) 4.7 x 10 ⁹ (>30 Mev)
17			2,100 (>20 Mev) (p/cm ² -sec-ster)	2.2 x 10 ⁸ (>20 Mev) p/cm ² -ster
18				2-3 x 10 ⁸ (>40 Mev) 4.8 - 7.3 x 10 ⁸ (>80 Mev) 2.6 - 4.5 x 10 ⁸ (>100 Mev)
19		10 (>30 Mev)	12,320 (>30 Mev)	
20	16 (>10 to 30 Mev)		5000 (>10 Mev) (p/cm ² -sec-ster) (21 db)	
21				1.6 x 10 ⁸ $\frac{p}{\text{cm}^2} > * \text{Mev}$
22				2.0 x 10 ⁸ $\frac{p}{\text{cm}^2} \frac{p}{\text{cm}^2\text{-ster}} 20 \text{ Mev}$
23			1-10 x 10 ³ $\frac{p}{\text{cm}^2\text{-sec-ster}}$	
24			50 $\frac{p}{\text{cm}^2\text{-sec-ster}} > 80 \text{ Mev}$	$\alpha E - 3 \pm 0.3$ at t = 30 hr
*Minimum energy not given.				



A good summary of measurements made on this event is presented by Masley and Goedeke (Reference 32). However, it is our opinion that the 4π isotropy assumed throughout their analysis for the rocket and balloon altitude flux measurements is incorrect, especially since it results in a proton flux approximately six times that reported by other authors. This problem illustrates the difficulty of obtaining integrated flux numbers for the various events, since they must be inferred from indirect measurements (neutron monitors, riometers, etc.) and from a few flux rate measurements.

Since the numbers reported by Webber (Reference 4) agree with those selected for integral fluxes, peak flux rates, rise and characteristic decay times >30 and >100 Mev, his values were used for the integral flux and peak flux rate >10 Mev as well. However, had reported values for these parameters been unavailable, the technique for estimating them is straightforward. Based upon the available numbers, a value of A would have been chosen (Figures 19-22). For the event of 12 November 1960, A is $\sim 1.0 \times 10^8$, which leads to $\int \phi > 10 \text{ Mev} = 9 \times 10^9 \text{ p/cm}^2$, and $\hat{\phi} > 10 \text{ Mev} = 36,000 \text{ p/cm}^2\text{-sec}$. These are the values that would have been used had not reliable published numbers ($\int \phi > 10 \text{ Mev} = 4 \times 10^9 \text{ p/cm}^2$ and $\hat{\phi} > 10 \text{ Mev} = 32,000 \text{ p/cm}^2\text{-sec}$) been available. This not only illustrates how mission data were estimated but given an idea of the accuracy of the estimation technique used.

A number of very small events reported by Krimigis, Van Allen, and Frank (Reference 33), Van Allen (Reference 34), and Cline et al (Reference 35) have not been included in the table because of their size. Undoubtedly, there have been hundreds of such small events which have not been detected. However, it is the large events ($>10^7 \text{ p/cm}^2$ >30 Mev) that control the annual proton fluxes and are of major interest from biological and material damage standpoints.



III. CORRELATIONS WITH SOLAR PARAMETERS

SELECTION OF CORRELATIONS

On the basis of the solar proton radiation event data presented in the preceding section, the future occurrence of events must be expected to be essentially random unless a means of predicting events can be established. It is clearly impossible to predict individual events farther into the future than the life-time of a typical active solar region, i. e., a few months. Since it is desired to evaluate the solar proton radiation environment in the 1969 to 1975 time period, an index of solar activity must be found which can be predicted several years in advance and which is strongly correlated with the mean flux of energetic protons associated with flares and with the frequency of solar proton radiation events.

Because of the generally cyclic but not fully periodic nature of solar activity, a reliable prediction of the future level of any activity index must be based on knowledge of past activity over many cycles. The sunspot number* is the only adequate activity index, having been observed continuously and uniformly since 1749, while other indices have been recorded only since 1913 or later. Sunspot data, therefore, exist for 19 complete cycles of mean duration 11.2 years, whereas no other activity data are consistently available for more than four cycles. As will be shown later, trends exist in the sunspot cycle which represent systematic changes over periods as long as 89 years; therefore, activity indices that have been recorded only for shorter periods are less capable of providing activity forecasts for future years.

The long-range prediction of mean levels of solar activity and flare-associated proton flux can be supplemented by methods applicable to the forecasting of individual proton radiation events from one to 35 days in advance. These methods are based on statistically significant relationships between observable characteristics of individual active regions and the future occurrence of solar flares at these regions. The utility of short-range forecasts of flares and, hence, of the likelihood of increases in the proton radiation environment level, for interplanetary missions lies in the ability to program and command evasive maneuvers (such as presentation of thick spacecraft sections toward the direction of approach of the most energetic protons) and the alteration of equipment operating modes (such as the telemetry cycle of a charged particle spectrometer). It is obvious that, in a flight

*The term "sunspot number" in this report refers to the monthly smoothed Wolf number (Zürich).



of several hundred days duration, at least one solar proton event will almost certainly occur. Short-range solar activity forecasts, therefore, are relevant to detailed spacecraft operations rather than to mission scheduling.

The characteristics of active solar regions selected as bases of flares prediction are the area, brightness, and luminosity of calcium plage regions and the presence of bipolar or complex magnetic regions in which the line separating areas of opposite polarity has no proper motion with respect to the solar photosphere (Reference 36).

Brightening in the chromospheric Ca II K₃ line is one of the earliest and most stable optical indications of the formation of an active region; therefore, plages promise a good opportunity for advance warning of flares. More transitory optical indices, such as the area of sunspot groups, do not afford sufficient stability for the reliable forecasting of activity more than a few (about three) days in advance (Reference 37).

Magnetic activity in a photospheric region often precedes optically observable activity and, as will be shown, allows the forecasting of flare occurrence probability with greater accuracy than is possible by means of plage observations alone.

An additional correlation, between solar proton events and previous slight decreases in the neutron flux near sea level, was investigated.

PROTON FLUX - SUNSPOT NUMBER

In order to evaluate the probability distribution of the mean flare-associated proton flux versus proton energy in 1969 to 1975, the correlation between this flux and the sunspot number must be established, and the probable sunspot number during these years must be estimated.

Sunspot Number Prediction

Although certain partial regularities, such as the basic cycle of mean duration 11.2 years, are evident in the temporal dependence of the sunspot number, the detailed behavior of this number is sufficiently complicated to require representation by a broad spectrum of component frequencies. If only the most important component frequencies such as those at 0.0112, 0.089, and 0.100 year⁻¹ are used, the fluctuations in the sunspot number are still inadequately represented. Although there are presumably physical reasons underlying the sunspot cycle and its correlation with solar flare proton radiation occurrences, these reasons are too poorly understood to permit a theoretical and analytical approach to the problem of forecasting solar proton flux occurrence probabilities. It is necessary, therefore, to adopt a



statistical approach which takes into account what may be truly random variations in the sunspot number and the solar flare proton flux level.

Let $R_M(t)$ be the monthly smoothed Wolf sunspot number at time t , where $t = 0$ on 1 July 1749 (Reference 38). The integral

$$\int_0^{t'} R_M(t) e^{-i 2\pi f t} dt = \int_0^{t'} R_M(t) (\cos 2\pi f t - i \sin 2\pi f t) dt \quad (2)$$

has the amplitude which, when averaged over an infinite time interval, is equivalent to the sum of the Fourier sine and cosine integrals of $R_M(t)$ (Reference 39),

$$\begin{aligned} P(f) &= \lim_{t' \rightarrow \infty} \frac{1}{t'} \left| \int_0^{t'} R_M(t) e^{-i 2\pi f t} dt \right|^2 \\ &= \left[\int_0^{\infty} R_M(t) \sin 2\pi f t dt \right]^2 \\ &\quad + \left[\int_0^{\infty} R_M(t) \cos 2\pi f t dt \right]^2 \end{aligned} \quad (3)$$

because of the orthogonality of the sine and cosine functions. The "power" P , i. e., the amplitude squared at frequency f , is defined as the frequency power spectrum of the sunspot number $R_M(t)$. Since R_M is known only during an interval of 216 years (actually a 215-year interval, 1749-1964, was used in this program), the limit in Equation 3 was approximated by an upper bound of $t' = 215$ years; therefore, the lower bound on f is $4.65 \times 10^{-3} \text{ year}^{-1}$. An upper bound, $f_0 = 0.2 \text{ year}^{-1}$, was chosen because of the essential randomness of rapid fluctuations in R_M . The resulting spectrum of $P(f)$ versus f is shown in Figure 1 (Reference 40), and the values and amplitudes of the 25 largest frequency components are presented in Table C2. Inspection of Figure 1 shows that any other components of the frequency power spectrum are negligible. Since $R_M(t'')$ is uniformly continuous piecewise on the interval $(1749.5 \leq t'' \leq 1976.0)$, the only uncertainty in the use of Equation 4 is the one arising from the use of a finite number of components of the power frequency spectrum (Reference 39).



The sunspot number at time t'' is the average of the frequency amplitude spectrum over the range of frequencies,

$$R_M(t'') = \left[\frac{1}{f_o} \int_{1/t'_m}^{f_o} P(f) \sin 2\pi f (t'' - t_o) df \right]^{1/2} \quad (4)$$

where $t'_m = 215$ years and t_o is a time at which $R_M(t_o) = 0$. This value occurred only from February through December 1810, and so $t_o = 61.5 \pm 0.5$ year (after 1 July 1749). Then $R_M(t'') = 0$ at $t'' = 1810.5 \pm 0.5$, in accord with observation.

$R_M(t'')$ was calculated at each month from July 1749 through December 1969 by means of the computer program SPOTNO described and listed in Appendix C. The results were normalized so that the arithmetic mean deviation of predicted and observed maxima between 1749 and 1964 was zero. The time scale was translated so that the arithmetic mean deviation of predicted and observed dates of maxima between 1749 and 1964 was zero; this procedure is equivalent to the choice $t_o = 65.2$ year. In the normalization procedure, the un-normalized predicted maxima of $R_M(t'')$ have a mean value $\overline{R_M'} = 49.2$, and the observed maxima of cycles 1 to 19 have a mean value $\overline{R_M} = 108.1$. The calculated values of $R_M(t'')$ are then multiplied by the factor $(\overline{R_M}/\overline{R_M'}) = 2.20$. In the translation procedure, the unadjusted predicted maxima occur at times t'_i , $i = 1$ to 19, and the observed maxima occur at times t_i . The translation adds to each t'_i the amount $\delta t = 0.47$ year, so that

$$\sum_{i=1}^{19} (t'_i + \delta t - t_i) = 0$$

The dates and values of observed and predicted maxima of R_M are presented in Table 3. Observed and predicted values of R_M are shown in Figure 2 for recent cycles (15 through 19), and predicted monthly values during cycle 20 (January 1969 through December 1975) are also given. The distribution of deviations of observed and predicted maximum values of R_M is plotted in Figure 3 as a function of the standard deviation $\sigma_R = 54.5$. This distribution is fitted by a Gaussian distribution within statistical errors resulting from the population (19) of the sample of deviation values. The deviation of predicted from observed values of R_M is, therefore, random and not systematic. Comparison of predicted and observed sunspot numbers in the early part of cycle 20 (1964 to date) (Figure 2), shows reasonable agreement.




Figure 1. Sunspot Number Frequency Spectrum



An analysis (Reference 41) of the frequency distribution versus amplitude of sunspot number maxima led to the prediction of a maximum number of 135 in 1968, with probability 0.75 that this maximum will lie in the range $110 < R_M < 160$.

Using a correlation of R_M maxima with rise times from minimum to maximum, King-Hele (Reference 42) has predicted a maximum of $R_M = 140$ in February 1968; he does not give the standard deviations of these values.

As an alternate approach to development of a formal representation of the behavior of the sunspot number, an attempt was made to fit $R_M(t)$ by a polynomial

$$R_M(t) = \sum_{n=0}^n a_n (t-t_0)^n \quad (5)$$

by a least-squares technique. Since there are 19 observed maxima, a polynomial of degree 19 or greater is required. The behavior of such a polynomial is so sensitive to the terms of higher degree at large values of $(t-t_0)$ that the terms of low degree can not feasibly be evaluated with sufficient precision for accurate determination of $R_M(t)$ near $t = t_0$.

Proton Flux - Sunspot Number Correlation

The results presented in Section II provide values of the annual proton flux $\int \phi_Y (>E)$ at proton energies greater than E , at solar distance 1 AU. Figure 4 shows the annual flux above 10, 30, and 100 Mev and the annual smoothed Wolf sunspot number R_Y for 1955 through 1964. Use of annual proton flux values reduces fluctuations associated with single large or anomalous solar proton events such as that of 23 February 1956, without destroying the general correlation of the flux with the sunspot number.

The correlation coefficient of the annual smoothed Wolf sunspot number with the common logarithm of the annual proton flux is

$$C = \frac{\sum_{i=1}^I (R_{Yi} - \bar{R}_Y) \left(\int \phi'_{Yi} - \overline{\int \phi'_Y} \right)}{\left[\sum_{i=1}^I (R_{Yi} - \bar{R}_Y)^2 \sum_{i=1}^I \left(\int \phi'_{Yi} - \overline{\int \phi'_Y} \right)^2 \right]^{1/2}} \quad (6)$$



Table 3. Monthly Smoothed Wolf Sunspot Numbers at Maxima

Observed			Predicted	
Cycle	Year	Number	Year	Number
1	1761.5	86.5	1759.4	70.1
2	1769.7	115.8	1770.4	90.6
3	1778.4	158.5	1781.4	71.2
4	1788.1	141.2	1791.7	16.5
5	1805.2	49.2	1802.2	74.4
6	1816.4	48.7	1812.1	110.6
7	1829.9	71.7	1827.3	140.0
8	1837.2	146.9	1837.8	165.3
9	1848.1	131.6	1847.9	140.0
10	1860.1	97.9	1859.1	110.6
11	1870.6	140.5	1869.0	74.4
12	1883.9	74.6	1884.3	16.5
13	1894.1	87.9	1894.7	71.2
14	1907.0	64.2	1905.7	71.6
15	1917.6	105.4	1916.2	99.5
16	1928.4	78.1	1927.4	91.0
17	1937.4	119.2	1937.5	115.0
18	1947.5	151.8	1947.4	105.4
19	1957.0	190.2	1955.8	103.4
20	--	--	1969.8 \pm 0.7	135.2 \pm 54.5




Figure 2. Observed and Predicted Sunspot Numbers




Figure 3. Distribution of Deviations of Predicted Maximum Sunspot Numbers



where I is the number of years sampled (10) and $\int \phi'_Y = \log_{10} \int \phi_Y$ was calculated for threshold energies of 10, 30, and 100 Mev. The results, presented in Table 4, are only slightly affected by omission of the 23 February 1956 event, the most unusual of the larger events in regard to its hard spectral distribution.

Table 4. Correlation of Proton Flux with Sunspot Number

Minimum Proton Energy	Correlation Coefficient
10 Mev	0.67
30 Mev	0.60
100 Mev	0.53

The ellipses in Figure 4 indicate the 95-percent confidence level that the flux in a given year from 1969 to 1975 will lie in the vertical range bounded by each ellipse at the abscissa corresponding to that year*. The most probable flux values lie on the major axis at these abscissae. These ellipses have equations of the form (Reference 43), adjusted for a sloping major axis,

$$\frac{R_Y^2}{\sigma_R^2} - \frac{2 CR_Y \int \phi'_Y}{\sigma_R \sigma_\phi} + \frac{(\int \phi'_Y)^2}{\sigma_\phi^2} = (0.95)^2 \quad (7)$$

where σ_R^2 is the sum of the variances of the sampled annual sunspot number R_Y (1956 - 1963) and of the predicted monthly numbers (Equation 4); σ_ϕ^2 is the variance of the annual proton flux. The mean values of R_Y and $\int \phi'_Y$ have not been subtracted from the R_Y and $\int \phi'_Y$ in computing σ_R^2 and σ_ϕ^2 . The application of these results to the problem of evaluating the expected proton flux in the 1969-1975 time period is discussed further below and in Section IV.

Since the major axis of each ellipse is the locus of the most probable values of the annual flux above the corresponding energy*, and the probability distribution of $\log_{10} \int \phi_Y = \int \phi'_Y$ is assumed symmetrical about the most probable value, the probability is 0.5 that $\int \phi'_Y$ lies above the major axis. If a factor $p^2 < 1$ is used in place of $(0.95)^2$ on the right-hand side of Equation 7, the resulting ellipse represents the confidence level p of $\int \phi'_Y$. As p^2 is varied in the range $0 < p^2 < 1$, a set of confocal ellipses with coincident major axes is generated.

* The upper and lower limits of the distribution are indicated by the ellipses.




Figure 4. Annual Proton Fluxes Versus Sunspot Number



PROTON EVENT NUMBER - SUNSPOT NUMBER

There were 76 events reported in the literature in sufficient detail to warrant their inclusion in Table 1. It was necessary to correlate these data with the monthly smoothed Wolf sunspot numbers if the predicted sunspot numbers for the next solar cycle (cycle 20) are to be used as a basis for predicting future proton events.

In order to seek a correlation between the proton event data and the sunspot number, it seemed logical to arrange the proton event data in a pattern which provided only a small amount of bias with event size. This may be accomplished by constructing Table 5 from the data in Table 1. One count is entered in each column to the left of the column corresponding to a flux greater than the flux in a given event. The integral fluxes above 30 Mev were used since these data were the most numerous and probably the most reliable as well. It will be noted that the intervals are open-ended, i. e., an event with $\int \phi > 10^9$ p/cm² above 30 Mev would be recorded as a 1 in each column. Hence, even though only two events were recorded for 1942, each had $\int \phi > 10^8$ p/cm², and so were recorded in the first three columns of Table 5.

Based upon Table 5, a proton event number (PEN), analogous to the annual smoothed sunspot number R_Y , may be defined as

$$PEN = \sum_i^N n_i \quad (8)$$

where n_i is the number of columns in Table 5 in which the event would be listed, and N is the number of actual observed events in the year. Values of N are also given in Table 5. For the purposes of this analysis, it is assumed that R_Y and the PEN are related thus

$$(PEN)_Y = 0.1 R_{Y-1} \pm 4.0 \quad (9)$$

Since the estimated R_Y for 1969 is 135 ± 54 , the estimated PEN for 1970 is 14 ± 5 . The proton event numbers PEN for the years $Y = 1940$ to 1964 and the sunspot numbers R_Y are plotted versus time in Figure 5. It is to be noticed that for cycle 19 (the only one for which shape comparisons are possible) there is a time lag of about one year, as indicated in Equation 9, between the sunspot number (which reached a maximum in 1957-1958) and the proton event number (which reached a maximum in 1959). Callender, Manzano, and Winckler (Reference 44) measured an 18-month time lag in cosmic ray ionization. Until more definitive data become available, it is not possible to assume that such a time lag does not exist.



Table 5. Proton Event Frequency Versus Flux Above 30 Mev

Year	Number of Observed Events	Number of Events with $\int \phi > 30 \text{ Mev Above Given Values (p/cm}^2)$					Total (PEN)
		$\int \phi > 10^7$	$\int \phi > 3 \times 10^7$	$\int \phi > 10^8$	$\int \phi > 3 \times 10^8$	$\int \phi > 10^9$	
1942	2	2	2	2	0	0	6
1946	1	1	1	1	0	0	3
1949	2	2	1	1	1	0	5
1951	2	2	2	0	0	0	4
1952	1	1	1	0	0	0	2
1956	3	3	2	2	1	1	9
1957	14	8	7	3	1	0	19
1958	11	6	6	4	1	0	17
1959	7	6	5	4	4	2	21
1960	17	4	4	2	2	1	13
1961	10	4	3	1	1	0	9
1962	2	0	0	0	0	0	0
1963	4	2	2	1	1	0	6
Total	76	41	36	21	12	4	114

The time lag of approximately one year was obtained by comparing the standard deviations for the PEN from 0.1 Ry for the following three cases.

Time Lag	Standard Deviation
0	4.2
1 year	4.0
2 years	7.4




Figure 5. Proton Event Number and Sunspot Number, 1940-1964



Knowing the proton event number is not sufficient to calculate the total proton flux expected in any future year. A total of 14 events with $\int \phi = 10^7$ $\text{p/cm}^2 > 30$ Mev would yield a PEN of 14 and a total flux of 1.4×10^8 $\text{p/cm}^2 > 30$ Mev. However, the same PEN would result if there were two events of 10^9 $\text{p/cm}^2 > 30$ Mev and one event of 5×10^8 $\text{p/cm}^2 > 30$ Mev, but the total proton flux would be almost 20 times greater. This, together with the use of the PEN concept, is illustrated in Section V.

FLARE-ACTIVE SOLAR REGION

While the correlation between proton flux and the sunspot number is most valuable to interplanetary spacecraft radiation protection analysis, the ability to predict the probability of occurrence of solar flares during specific periods is also useful. Justification of the selection of characteristics of plage regions and magnetically active regions as indicators of future flare occurrence lies in the expectation that a strong correlation of flares with these indices will exist in future years. In order to evaluate the flare-plage correlation, let the plage areas, ages (numbers of disk passages), and luminosities as observed at central meridian passage (CMP) be assigned to intervals with indices i , j , k , respectively, defined in Table 6.

It must be emphasized that the correlations discussed here permit only the prediction of solar flare occurrence probabilities and not the occurrence or size of proton radiation events. A high probability of flare occurrence, however, is associated with a high probability of occurrence of enhanced levels of energetic proton radiation.

Table 6. Intervals of Parameters of Plage Regions

i	Area ¹	j	Age ²	k	Luminosity ³
1	<1000	1	1	1	1.0, 1.5
2	1000-2000	2	2	2	2.0, 2.5
3	2000-4000	3	3	3	3.0, 3.5
4	≥ 4000	4	4	4	4.0 - 5.0
		5	5		
		6	6		
		7	7		
		8	≥ 8		
¹ Millionths of solar hemisphere. ² Solar rotations. ³ Units on McMath-Hulbert scale.					



Let P_{ijk} be the number of plages in one year which had area in the i th interval, age in the j th interval, and luminosity in the k th interval. For example, P_{243} is the number of plages with area from 1000 to 1999 millionths of the hemisphere which were appearing on the visible hemisphere for the fourth time and had luminosity or intensity 3 or 3.5 on the McMath-Hulbert observatory scale. Let P'_{ijk} be the number of plages with similar characteristics, at each of which one or more flares occurred during the disk passage following the passage on which the given characteristics were observed. The flare-plage associations are those contained in Reference 45. In computing P'_{ijk} , only flares with corrected area 100 millionths of the solar hemisphere or greater are considered. Smaller flares have poorer correlation with plage characteristics and are not predictable with significant accuracy by the present method. If consideration were limited to flares larger than, for example, 200 millionths of the hemisphere, the smallness of the data sample would reduce statistical accuracy. The term "flare" will be understood in this section as one with corrected area at least 100 millionths of the hemisphere.

Let w_{ijk} be the probability that at least one flare will occur at any given plage region, during the disk passage following the one on which the CMP area, age, and luminosity of the plage were in intervals i , j , and k , respectively. A good estimate of w_{ijk} is given by (Reference 36)

$$w_{ijk} = P'_{ijk} / P_{ijk} \quad (10)$$

If the characteristics of a plage are observed at CMP, then w_{ijk} is known at least 21 days before the occurrence of any flare considered in association with this plage. If the region is formed west of the central meridian, this time is 14 days. The values of w_{ijk} for each combination of plage characteristics are listed in Appendix B for the years 1958 to 1964; these are assumed applicable to the years 1969 to 1975, respectively, in the next solar activity cycle.

Similarly, let P''_{ijk} be the number of plages in area interval i , age interval j , and luminosity interval k at which one or more flares occurred during the disk passage at which the plage was observed. Also let

$$w''_{ijk} = P''_{ijk} / P_{ijk} \quad (11)$$

Values of w''_{ijk} , tabulated in Appendix B, represent the probability of flare occurrence within seven days before or after the plage CMP date. If the plage is observed, for example, at the solar equator at 60 degrees east longitude, w''_{ijk} approximates the probability of flare occurrence at this plage during the remaining 11.5 days of its disk passage.



Expected Accuracy of Predictions

If a flare occurs at a given region, the prediction (expressed as a probability) is considered to have a fractional correctness of w_{ijk} . If no flare occurs at this region, the prediction has a fractional correctness of $(1-w_{ijk})$. In a given time interval, $\sum_{ijk} P_{ijk}$ predictions are made, one for each plage region.

The notation $\sum_{ijk} P_{ijk}$ represents the triple summation $\sum_{i=1}^I \sum_{j=1}^J \sum_{k=1}^K P_{ijk}$ over the i , j , and k intervals to which the plage area, age, and luminosity, respectively, can be assigned. Reference to Table 6 shows that $i = 4$, $j = 8$, $k = 4$ in the present analysis. The expectation that any prediction will prove to be "correct" in the way correctness has been defined above is the weighted average of the fractional correctness of all regions (Reference 36).

$$S = \frac{\sum_{ijk} P'_{ijk} w_{ijk}}{\sum_{ijk} P_{ijk}} + \frac{\sum_{ijk} (P_{ijk} - P'_{ijk}) (1 - w_{ijk})}{\sum_{ijk} P_{ijk}}$$

$$S = \frac{\sum_{ijk} P_{ijk} (1 - 2 w_{ijk} + 2 w_{ijk}^2)}{\sum_{ijk} P_{ijk}} \quad (12)$$

The expectation that a "false alarm" will result, i. e., that no flare will occur at the given region is

$$S_F = \frac{\sum_{ijk} [(P_{ijk} - P'_{ijk}) w_{ijk}]}{\sum_{ijk} P_{ijk}} \quad (13)$$

$$S_F = \frac{1}{2} (1 - S) \quad (14)$$



If a prediction is neither correct nor a "false alarm," it represents a failure to predict a flare that does occur. The expectation of such a failure is

$$S_o = 1 - S - S_F = S_F \quad (15)$$

The expected accuracy of predictions of flare occurrence on the same disk passage as that at which the plage characteristics were observed can be represented by S'' , S_F'' and S_o'' , which are found by replacing P'_{ijk} with P''_{ijk} and w_{ijk} with w''_{ijk} in Equations 12, 14 and 15, respectively. Values of S and S'' are presented in Table 7 for the years 1958 through 1964. These values are assumed to hold for the corresponding years from 1969 through 1975, following an 11-year solar activity cycle. If the variables were uncorrelated, S or S'' would equal 0.5. The values of S apply only to flares at plages observable on the disk passage preceding the flare.

Table 7. Flare Prediction Accuracy

Year	Same Disk Passage (S'')	Next Disk Passage (S)
1958	0.716	0.674
1959	0.695	0.673
1960	0.713	0.888
1961	0.733	0.744
1962	0.717	0.717
1963	0.898	0.925
1964	0.899	0.960

Flare-Magnetic Field Correlation

Previous studies (References 36, 46) have established conditions on the solar magnetic field component normal to the photosphere, B_p , which are favorable to the occurrence of solar flares during the next disk passage of a magnetically active region. These simultaneous conditions are:

1. $|B_p| \geq 2$ gauss for 3 days or longer
2. $|\partial B_p / \partial t| \leq 1$ gauss/day
3. $|\nabla B_p| \geq 10$ gauss/heliographic degree



The gradient of B_p is understood to be the component normal to the line of sight of $\nabla (\vec{B} \cdot \vec{n})$ where \vec{n} is a vector of unit length parallel to the line of sight. This association is most useful in cases where magnetic activity is evident before the formation of a calcium plage at a new region, and permits forecasting of flare occurrence probability at dates 21 to 35 days in advance at some regions at which the plage is "new" at the disk passage on which the flares occur.

To evaluate this correlation, the positions of flares at new calcium plages were translated to the dates at which magnetograms were obtained during the preceding solar rotation. The translation included effects of differential solar rotation and the inclination of the solar equator to the ecliptic plane. Table 8 presents the number and fraction of plages with age 1 rotation (i. e., new plages) in each year from 1960 through 1964; these plages

Table 8. Magnetic Field—Flare Association

Year	New Plages		Stationary Magnetic Null Points		Other New Flaring Regions
	Number	Fraction	Total	Flaring	
1960	123	0.291	90	72	23
1961	101	0.343	38	20	21
1962	162	0.535	22	14	15
1963-4	883	0.917	12	4	11

are the ones to which the magnetic field-flare correlation was applied. The table gives the number of cases in which the magnetic field conditions listed above occurred at the regions at which new plages were to appear on the solar rotation following the magnetic observation. Next is given the number of cases in which the existence of these magnetic conditions was followed by one or more flares on the next rotation. Finally the table indicates the number of new and flaring plage regions at which the magnetic conditions did not occur. Magnetograms prepared before 1960 (Reference 47) lack sufficient angular resolution for use in this analysis. The years 1963 and 1964, near a minimum stage of solar activity, were combined to provide a more significant statistical sample.



Accuracy of Combined Plage and Magnetic Correlations

Let N_M be the number of stationary magnetic null points in one year, i. e., the number of cases in which the previously described magnetic conditions occurred. Let N'_M be the number of such points near which flares were observed on the following solar rotation*. The combined accuracy of flare occurrence predictions based on association with plages and with magnetic field patterns is defined as

$$S^* = S (1 - \sum_{ik} P_{ilk}) / \sum_{ijk} P_{ijk} + N'_M \sum_{ik} P_{ilk} / N_M \sum_{ijk} P_{ijk} \quad (16)$$

where S is defined in Equation 12. The sum $\sum_{ik} P_{ilk}$ is the number of new plages ($j = 1$) in the yearly data sample. Values of S^* are given in Table 9; the results in Tables 8 and 9 are assumed applicable to the years 1971 through 1975. In the absence of results for 1958 and 1959, it is assumed that the results for 1960 apply to 1969 and 1970.

The results shown in Table 9 indicate that flare occurrence probabilities can be predicted with reasonable confidence during years of high solar activity, but with only very limited validity during quiet years. The usefulness of this prediction technique in interplanetary radiation environment analysis has been discussed at the beginning of Section III.

Table 9. Accuracy of Combined Flare Prediction Method

Year	S^*
1960	0.862
1961	0.590
1962	0.674
1963	0.38
1964	0.38

* "Near" is defined as within 10 heliographic degrees.



PROTON EVENT-NEUTRON MONITOR RATE

From one to fourteen days before some polar cap absorption (PCA) events, decreases of one to three percent have been observed in the neutron flux near sea level (References 48, 49). These so-called pre-decreases (pré-baisses) are to be distinguished from the larger Forbush decreases of typically 10 to 20 percent which accompany rather than precede some solar energetic proton events. Pre-decreases are probably caused by the exclusion from the earth of galactic charged particles by a plasma concentration that is associated with an active solar region and that lies to one side of the earth. Legrand's tabulation of pre-decreases from 26 February 1957 to 18 December 1957, when compared with the proton event list of Table 1, reveals the following association:

Total pre-decreases	12
Pre-decreases followed within 14 days by PCA event	6
Pre-decreases followed by geomagnetic storm but not PCA event	2
PCA events not preceded by decrease	2

The probability of a successful prediction of a polar cap absorption event is accordingly 0.50 ± 0.17 , with 75 percent of proton events falling within the alert period from one to fourteen days after a pre-decrease. The validity of this method is rather limited, but it provides warnings during periods inaccessible by optical techniques because of the 14-day period required for passage of an active region across the solar disk.



IV. PROTON RADIATION OCCURRENCE PROBABILITY

MODEL EVENT AT 1 AU

In order to obtain a model solar proton event at 1 AU from the sun, it was necessary to establish correlations in the data that were available. The available data consisted of onset plus rise times for protons above 30 and above 100 Mev, characteristic decay times for the same protons, and peak flux rates and integral fluxes for protons above 10, 30, and 100 Mev. These data are listed in Table 1. It was these data that were examined for patterns that would permit the construction of a model event at 1 AU.

A previous study (Reference 50) has been carried out along these lines, which involved forming the following ratios which are subject to estimated standard deviations as shown:

$$\frac{\sum \int \phi > 10 \text{ Mev}}{\sum \int \phi > 30 \text{ Mev}} = 5.14 \pm 0.27 \sim E^{-1.51} \quad (17)$$

$$\frac{\sum \int \phi > 30 \text{ Mev}}{\sum \int \phi > 100 \text{ Mev}} = 6.55 \pm 0.65 \sim E^{-1.57} \quad (18)$$

$$\frac{\sum \hat{\phi} > 10 \text{ Mev}}{\sum \hat{\phi} > 30 \text{ Mev}} = 3.4 \pm 0.3 \sim E^{-1.12} \quad (19)$$

$$\frac{\sum \hat{\phi} > 30 \text{ Mev}}{\sum \hat{\phi} > 100 \text{ Mev}} = 4.0 \pm 0.4 \sim E^{-1.16} \quad (20)$$

In the above expressions, $\int \phi$ is the total proton flux (p/cm^2) above the indicated energy and $\hat{\phi}$ is the peak flux rate ($p/cm^2\text{-sec}$). The summations were carried out in each case over all events for which the data required in both the numerators and denominators were available. In other words, in the first ratio, only those events were included for which both the integral fluxes above 10 Mev and the integral fluxes above 30 Mev were available. From these ratios, it was determined that the average integral energy spectrum weighted by flux per event varied as $E^{-1.55 \pm 0.10}$, and the weighted peak flux rate varied as $E^{-1.15 \pm 0.09}$ (see Reference 50).



At this point, recourse was made to a mathematical expression used to fit the Bailey Model Event (References 2, 51). That expression was of the form

$$\phi(E > E_o, t) = \frac{A t e^{-\alpha E_o^n t}}{E_o^m} \quad (21)$$

where t = time after the flare (hours) and E = energy (Mev). A , α , m , and n were taken as constants. It is seen that the time integral of this expression from $t = 0$ to $t = \infty$ is:

$$\int \phi(E > E_o) = \frac{A}{\alpha^2 E_o^{m+2n}} \sim \frac{1}{E_o^{1.55 \pm 0.10}} \quad (22)$$

The peak flux rate, obtained when $t = \frac{1}{\alpha E_o^n}$, is

$$\hat{\phi}(E > E_o) = \frac{A e^{-1}}{\alpha E_o^{m+n}} \sim \frac{1}{E_o^{1.15 \pm 0.09}} \quad (23)$$

Thus, the first four ratios (Equations 17 through 20) formed from the data can be approximately fit, providing $m = 0.75 \pm 0.15$ and $n = 0.4 \pm 0.1$. The parameter α was evaluated by setting the time interval from $t = 0$ to $t = (1/\alpha E^n)$ equal to the size-weighted onset plus rise time. These onset plus rise times are shown in Figures 6 and 7 and were averaged, obtaining 12 ± 7 hours (for protons above 30 Mev) and 7 ± 5 hours (for protons above 100 Mev). Thus, $\alpha = 0.022 \pm 0.002$ and the rise time does vary approximately as $E^{-0.4}$.

The characteristic decay time as given by Equation 21 is not a constant, since the linear factor of t tends to offset the exponential. If, however, the time required for the peak flux rate to decrease a factor of e is used, the characteristic decay time obtained is ~ 2.15 times the onset plus rise time. This yields characteristic decay times of $\sim 25 \pm 12$ hours (for protons above 30 Mev) and $\sim 15 \pm 2$ hours (for protons above 100 Mev). As Figures 8 and 9 show, these are in reasonable agreement with the size-weighted data. It will be noted that t and τ are plotted to the nearest hour.

The following expressions may be derived from Equation 21.

$$\phi(E > E_o, t) = \frac{A t e^{-0.022 E_o^{0.4} t}}{E_o^{0.75}} \quad (\text{p/cm}^2\text{-hr}) \quad (24)$$




Figure 6. Onset Plus Rise Times Weighted for Flux Above 30 Mev





Figure 7. Onset Plus Rise Times Weighted for Flux Above 100 Mev



τ CHARACTERISTIC DECAY TIME (HRS)

Figure 8. Characteristic Decay Times Weighted for Flux Above 30 Mev




Figure 9. Characteristic Decay Times Weighted for Flux Above 100 Mev



$$\phi(E, t) = \frac{At e^{-0.022E^{0.4}t}}{E^{1.75}} (0.75 + 0.0088 t E^{0.4}) \text{ (p/cm}^2\text{-hr-Mev)} \quad (25)$$

$$\phi(E) = \frac{3200 A}{E^{2.55}} \text{ (p/cm}^2\text{-Mev)} \quad (26)$$

$$\int \phi(E > E_o) = \frac{2100A}{E_o^{1.55}} \text{ (p/cm}^2\text{)} \quad (27)$$

$$\hat{\phi}(E > E_o) = \frac{17A}{E_o^{1.15}} \text{ (p/cm}^2\text{-hr)} \quad (28)$$

$$t_{\text{rise}}(E > E_o) = \frac{45}{E_o^{0.4}} \text{ (hr)} \quad (29)$$

$$\tau_{\text{decay}}(E > E_o) = \frac{100}{E_o^{0.4}} \text{ (hr)} \quad (30)$$

where A is a normalization constant determined from $\int \phi$ by Equation 27.

These expressions were derived as part of a series of radiation shielding studies carried out previously (Reference 50). Their use in the present study was hampered by their applicability to only the largest events. In order to obtain a model applicable to events of any size, modifications were required. In order to determine the sort of modifications required, the data were examined for characteristics that could be correlated with event size. Two trends were noted. The slopes of the integral energy spectra (Figure 10) and the peak flux spectra (Figure 11) tend to become less steep, and the ratio of integral flux to peak flux rate increases (Figures 12, 13, and 14) as the event size increases. Specifically, the integral energy spectrum appears to vary from $\sim E^{-1.6}$ for large events ($\int \phi > 30$ Mev above 10^9 p/cm²) to $\sim E^{-2.6}$ for small events ($\int \phi > 30$ Mev below 10^8 p/cm²). The relationship of integral flux to peak flux rate appears to be approximately (for all energies)

$$\int \phi \sim 2.5 \times 10^4 \hat{\phi}^{1.177} \quad (31)$$




Figure 10. Integral Fluxes of Several Proton Events




Figure 11. Peak Flux Rates of Several Proton Events




Figure 12. Integral Flux Versus Peak Flux Rate Above 10 Mev




Figure 13. Integral Flux Versus Peak Flux Rate Above 30 Mev




Figure 14. Integral Flux Versus Peak Flux Rate Above 100 Mev



In order to incorporate these trends into Equation 21, it was necessary to make α and m functions of event size (determined by the normalization factor A). The desired behavior can be approximated by the following relationships:

$$\alpha = A^{-0.224 \pm 0.01} \quad (32)$$

(from Equation 31) and

$$m = 16A^{-0.16 \pm 0.02} \quad (33)$$

which accounts for the $E_o^{-1.6}$ to $E_o^{-2.6}$ behavior. Substituting these into Equations 24 to 30 yields

$$\phi(E > E_o, t) = \frac{At e^{-\left(\frac{E_o^{0.4} t}{A^{0.224}}\right)}}{E_o^{16A^{-0.16}}} \text{ (p/cm}^2\text{-hr)} \quad (34)$$

$$\phi(E, t) = \frac{At e^{-\left(\frac{E^{0.4} t}{A^{0.224}}\right)}}{E^{16A^{-0.16} + 1} (0.4 t A^{-0.224} E^{0.4} + 16A^{-0.16})} \text{ (p/cm}^2\text{-hr-Mev)} \quad (35)$$

$$\int \phi(E > E_o) = \frac{A^{1.45}}{E_o^{16A^{-0.16} + 0.8}} \text{ (p/cm}^2\text{)} \quad (36)$$

$$\phi(E) = \frac{16A^{1.29} + 0.8A^{1.45}}{E^{16A^{-0.16} + 1.8}} \text{ (p/cm}^2\text{-Mev)} \quad (37)$$

$$\hat{\phi}(E > E_o) = \frac{0.37 A^{1.224}}{E_o^{16A^{-0.16} + 0.4}} \text{ (p/cm}^2\text{-hr)} \quad (38)$$

$$\tau_{\text{decay}}(E > E_o) = 2.15 t_{\text{rise}}(E > E_o) = \frac{2.15 A^{0.224}}{E_o^{0.4}} \text{ (hr)} \quad (39)$$



The solid curves in Figures 12, 13, and 14 were calculated using these expressions. It is seen that they fit the data fairly well.

As an additional check on Equation 39, the rise (actually onset plus rise) and characteristic decay times as a function of the integral flux were compared with the formula. This is essentially the same as comparing the ratio of integral flux to peak flux rate, but the data exhibit more scatter (see Figures 15 through 18). Nevertheless, the calculated curves fit the data about as well as any other set of single-valued curves.

In order to facilitate the use of this model in estimating the missing event data, a series of graphs was made in which various parameters (peak flux rate, integral flux, rise and decay times) were plotted as functions of the normalization factor A . These graphs are shown in Figures 19 through 22 and were quite useful since A is not linearly proportional to any single event parameter (see Equations 34 through 39).

By the use of Equations 34 through 39 (Figures 19 through 22), the missing data in Table 1 were estimated. These estimated numbers are indicated by asterisks (*) and were not plotted as data points in Figures 6 through 18.

The advantage of using an analytical model (Equations 34 through 39) rather than a numerical fit to the existing data (Figures 10 through 18) is that it is possible to generate all the characteristics of a solar proton event in a consistent pattern once a single parameter (e.g., t , τ , $\int \phi$ or $\hat{\phi}$) is known.

EVENT FREQUENCY DISTRIBUTION

As expected, ~~both the number and size of proton events are functions of~~ solar activity. The correlations of ~~event flux and frequency~~ with the sunspot number have been discussed in the previous section. In order to select events whose total flux ($\Sigma \int \phi$) and proton event number (PEN) add up to the totals expected on the basis of sunspot correlations, it is necessary to have an event frequency distribution to choose from. The most obvious distribution is by event size. In Figure 23, the number of proton events is shown as a function of integral flux above 10, 30, and 100 Mev. The 76 events of Table 1 were grouped into categories of $>10^{10}$ p/cm², $3 - 10 \times 10^9$ p/cm², $1 - 3 \times 10^9$ p/cm², $3 - 10 \times 10^8$ p/cm², etc. It is seen that the distributions are not in disagreement with the Gaussian distributions indicated by broken curves in Figure 23; the 1σ widths and means of the Gaussian curves are also shown in this figure. The lack of events larger than those observed is due presumably to their absence, while the lack of events smaller than those observed is due to observational difficulties. It is expected that if the proton event size distributions were completely known, they would be monotonic in nature, increasing as the event size decreased. However, these missed small events will not affect the annual proton fluxes appreciably.




Figure 15. Integral Flux Above 30 Mev Versus Onset Plus Rise Time




Figure 16. Integral Flux Above 30 Mev Versus Characteristic Decay Time




Figure 17. Integral Flux Above 100 Mev Versus Onset Plus Rise Time




Figure 18. Integral Flux Above 100 Mev Versus Characteristic Decay Time




Figure 19. Calculated Peak Flux Rates Versus Parameter A




Figure 20. Calculated Integral Flux Versus Parameter A





Figure 21. Calculated Onset Plus Rise Time Versus Parameter A



A

Figure 22. Calculated Characteristic Decay Time Versus Parameter A






Figure 23. Distributions of Integral Flux Per Event



The most probable size of an observed event is:

$$\int \phi > 10 \text{ Mev} = \sim 1.5 \times 10^8 \text{ p/cm}^2$$

$$\int \phi > 30 \text{ Mev} = \sim 2 \times 10^7 \text{ p/cm}^2$$

$$\int \phi > 100 \text{ Mev} = \sim 1.7 \times 10^6 \text{ p/cm}^2$$

These numbers have estimated 1σ limits of a factor of ± 2 .

A second type of event frequency distribution concerns the number of events per month. The 72 events listed in Table 1 represent a time interval of 22 years (~ 264 months) (1942 through 1963), but to conclude that there has been an average of 72/264 events per month would be erroneous because many of the early events undoubtedly have been missed. From 1956 on, the probability of missing an event $\gtrsim 10^4 \text{ p/cm}^2\text{-sec} > 30 \text{ Mev}$ has been fairly small, and 68 such events have been observed. Proceeding in this manner, the monthly probabilities of events as a function of event size over the 96 months from 1956 through 1963 can be obtained. The results are listed in Table 10. The differential percentages are listed at the left side of each column, the integral percentages are listed at the right. The maximum possible number in any column is 71 percent (68 events/96 months), and it is seen that the 50-percent probability/month event is rather small ($\sim 5 \times 10^6 \text{ p/cm}^2 > 30 \text{ Mev}$). The integral monthly probabilities (for events with $\int \phi$ above given values) are plotted in Figure 24.

It must be remembered that the monthly probabilities are averaged over the years 1956 through 1963. If the entire solar cycle 19 were included (~ 130 months), the monthly probabilities would probably have been decreased by ~ 25 percent, since the missing portions are near the solar minima. For the peak years of 1958-1960, the monthly probabilities would be increased. However, the limited data available do not permit this sort of analysis with much confidence.

The proton events are not distributed uniformly throughout the months, but they tend to occur in certain months. The monthly distributions for all 76 events listed in Table 1 are given in Table 11. The events missed prior to 1956 will not tend to bias the entire sample, as they would have the monthly probabilities computed above. The same data, expressed in percentages of the totals, are shown in Figure 25. It is seen that July has been by far the most active month, with 18.5 percent of the events, 24.5 percent of the proton event number (PEN), and 33 percent of the integral flux ($\int \phi$) above 30 Mev. December has apparently been the least active month, with 0 percent in all categories. However, this distribution must be viewed with



Table 10. Monthly Probabilities of Event Size for 1956-1693

Event Size (p/cm ²)	$\int \phi > 10 \text{ Mev}$ (Events/Month)	$\int \phi > 30 \text{ Mev}$ (Events/Month)	$\int \phi > 100 \text{ Mev}$ (Events/Month)
$> 1 \times 10^{10}$	0	0	0
$3 - 10 \times 10^9$	7.3	0	0
$> 3 \times 10^9$	7.3	0	0
$1 - 3 \times 10^9$	10.4	4.2	0
$> 1 \times 10^9$	17.7	4.2	0
$3 - 10 \times 10^8$	9.4	7.3	1.0
$> 3 \times 10^8$	27.1	10.5	1.0
$1 - 3 \times 10^8$	6.3	6.3	5.2
$> 1 \times 10^8$	33.4	16.8	6.2
$3 - 10 \times 10^7$	16.7	12.5	4.2
$> 3 \times 10^7$	50.1	29.3	10.4
$1 - 3 \times 10^7$	11.5	4.2	5.2
$> 1 \times 10^7$	61.6	33.5	15.6
$3 - 10 \times 10^6$	6.3	21.8	11.5
$> 3 \times 10^6$	67.9	55.3	27.1
$1 - 3 \times 10^6$	0	5.2	9.4
$> 1 \times 10^6$	67.9	60.5	36.5
$3 - 10 \times 10^5$	3.1	6.3	18.7
$> 3 \times 10^5$	71.0	66.8	55.2
$1 - 3 \times 10^5$	0	1.0	5.2
$> 1 \times 10^5$	71.0	69.8	60.4




Figure 24. Monthly Proton Event Occurrence Probability




Figure 25. Monthly Distribution of Events, Event Number, and Integral Flux



Table 11. Distribution of Events by Month, 1942-1963

Month	Number of Observed Events	Proton Event Number	$\int \phi > E \text{ (p/cm}^2\text{)}$		
			10 Mev	30 Mev	100 Mev
January	4	8	5.5×10^9	3.6×10^8	1.6×10^7
February	5	8	2.8×10^9	1.2×10^9	3.7×10^8
March	5	10	7.5×10^9	1.1×10^9	1.0×10^8
April	7	2	5.6×10^8	7.3×10^7	6.7×10^6
May	6	7	6.2×10^9	1.0×10^9	9.4×10^7
June	3	5	2.0×10^9	2.4×10^8	2.2×10^7
July	14	28	2.0×10^{10}	4.0×10^9	4.1×10^8
August	10	13	3.7×10^9	4.5×10^8	2.3×10^7
September	13	13	3.7×10^9	5.5×10^8	6.5×10^7
October	2	2	2×10^8	5×10^7	1×10^7
November	7	18	1.1×10^{10}	2.6×10^9	4.4×10^8
December	0	0	0	0	0
Totals	76	114	6.5×10^{10}	1.2×10^{10}	1.5×10^9

some suspicion. While the earth is 7 degrees above the solar equator in July, and most of the events in solar cycle 19 took place at northern solar latitudes, these two facts may very well be unrelated. An alternate (and probably equally plausible) explanation is that balloon measurements of solar proton events require high geomagnetic latitudes, which are far less attractive in December than in July in the Arctic.

One can conclude that the probability of a proton event varies inversely with some function (approximately the logarithm) of event size, and that both the number and size of the probable events increase when the sun is active. The monthly event probabilities (Figure 24) should be corrected for the appropriate portions of the solar cycle and perhaps for the actual months involved (Figure 25) before applying them to mission analysis studies.



DEPENDENCE ON SOLAR DISTANCE

It is sufficient for purposes of establishing radiation protection requirements to determine the dependence on solar distance of the total flux of flare-associated protons in each solar proton event, as a function of proton energy. The motion of a spacecraft during a proton radiation event has negligible effect on the radiation environment. It is obvious from the requirement of conservation of particles that the total flux at any energy should vary as the inverse square of the solar distance, unless the protons are confined to some channel or tube by interplanetary magnetic fields. A proof is presented later that this inverse-square dependence holds if the interplanetary magnetic field is that due to the solar field in the steadily expanding corona.

The adequate design of spacecraft instrumentation to measure the particulate radiation environment requires knowledge of the expected effect of solar distance on the temporal and angular dependence of the flare-associated proton flux as a function of proton energy. Consideration of angular dependence is beyond the scope of this program, although the evaluation of temporal dependence included analysis of the angular distribution of the proton flux versus energy and solar distance.

Theoretical Basis of Model

In order to express the dependence of the proton flux on energy, time, and solar distance, a model must be used which quantitatively describes the processes by which a solar flare transfers energy to protons in the solar atmosphere and by which the protons are transported through the interplanetary medium. The model selected in this work is based on:

1. Acceleration of protons by Lorentz forces in a hydromagnetic shock wave induced in the solar corona by absorption of electromagnetic radiation emitted by a flare
2. In the absence of local irregularities in the interplanetary medium, circular motion of energetic protons about guiding centers that move parallel to the interplanetary magnetic field vector
3. Random walk of energetic protons which enter local plasma concentrations containing disordered magnetic fields.

The results of these assumptions agree in many respects with the temporal and angular distribution of flare-associated proton energy spectra observed in the vicinity of the earth, as will be demonstrated.



Reasonable confidence, therefore, may be placed in the solar distance relationships based on this model.

Proton Acceleration by Hydromagnetic Shocks

It has been shown by Parker (Reference 52) that the sudden expansion of the solar corona, as a result of the absorption of electromagnetic radiation emitted by a flare, creates a hydromagnetic shock wave in the corona. This shock carries with it a disturbance of the interplanetary magnetic field, which does work on the solar wind protons overtaken by the shock. The accelerated protons are scattered forward of the shock by an approximately thermal process and are transported through the generally steady interplanetary magnetic field. The emergent protons remove energy from the shock wave, which gradually weakens and disappears. Superposition of the proton energy distributions emerging from shocks in all regions of the corona heated by the flare permits calculation of the time-dependent angular distribution versus energy of the protons arriving at a point at a given distance from the sun (Reference 54).

The magnetic field in the steadily expanding solar corona ahead of the shock is swept back azimuthally from the radial direction due to solar rotation relative to the coronal plasma. The components of the field in the solar equatorial plane are given by Parker (Reference 55) as

$$B_r(r, \theta, \phi) = B_r(r_1, \theta, \phi - r\Omega/u_o^x) (r_1/r)^2 \quad (40)$$

$$B_\theta(r, \theta, \phi) = 0 \quad (41)$$

$$B_\phi(r, \theta, \phi) = B_r(r_1, \theta, \phi - r\Omega/u_o^x) \left(\frac{r_1\Omega}{u_o^x} \right) \left(\frac{r_1}{r} \right) \sin \theta \quad (42)$$

where r , θ , and ϕ are the radial, polar, and azimuthal heliocentric coordinates; Ω is the sidereal angular velocity of the sun; and u_o^x is the solar wind transport velocity; i.e., the velocity of steady coronal expansion. When the leading edge of the shock wave is at $r = R_1$ and its trailing edge is at $r = R_2$, the magnetic field just inside the shock at $r > R_2$ is

$$B_{x_2} = B_{x_1} \quad (43)$$



$$B_{y_2} = B_{y_1} \left(\frac{f^2}{1-g} \right) \left(\frac{R_2}{R_1} \right)^4 \left(\frac{r}{R_2} - 1 \right)^{1-g} \quad (44)$$

$$B_{z_2} = B_{z_1} = 0 \quad (45)$$

In Equation 44, f and g are functions of the dimensionless shock parameter λ (Reference 52). A convenient form of the field inside the shock, reasonably accurate except when r is very slightly larger than R_2 , is the plane wave

$$B_i(x, y, z, t) = B_{i_1} + \left(\frac{x - vt}{x_2} \right) (B_{i_2} - B_{i_1}), \quad (46)$$

$$0 < t < t_2$$

$$x_2 = vt_2 \quad (47)$$

where $i = 1, 2, 3$ denotes the x -, y -, z -components, respectively, v is the velocity of the leading edge of the shock which passes the observer at $t = 0$, and the trailing edge passes him at $t = t_2$ (Reference 53).

If the magnetic field given by Equation 46, and a weak radial electrostatic field corresponding to a potential of 10^4 volt (the upper limit of solar wind proton energies is near 10^4 ev) between the sun and a point at 1 AU, are used in solution of the relativistic ponderomotive equations (References 53 and 54) of protons encountering the shock, the protons are found to reach energies from 1 Mev to about 400 Mev. Acceleration of a given proton terminates when it is scattered into the region in front of the shock. Coulomb and Rutherford scattering in the hot (5×10^5 to 2×10^6 degrees K) and nearly collisionless plasma are essentially random processes (Reference 54), so that the protons emerging from the shock have an isotropic angular distribution about the normal to the shock. The flux per unit time duration of the solar flare and the mean energy of the emerging protons are plotted in Figure 26 as functions of the heliocentric distance of the point of emergence. These results are for $\lambda = 4/3$ and are very similar to results for other values of λ in the allowed range $1 < \lambda < 3/2$.

Proton Transport in Regular Interplanetary Medium

A proton can reach a receiver point if the guiding center of its trajectory passes the receiver at a distance equal to the gyroradius measured at the receiver. The proton must emerge from the shock at a distance from the guiding center equal to the gyroradius at the point of emergence.




Figure 26. Proton Source Distribution Due to Hydromagnetic Shock



A typical trajectory is illustrated in Figure 27. It is a helical curve wrapped on a horn-shaped solid with a curving axis and an expanding cross-sectional radius.

In the unshocked corona, a magnetic line of force makes an angle with a heliocentric radial line given by (Reference 55)

$$\xi = \tan^{-1} (B_{\phi} / B_r) \quad (48)$$

$$\xi = \tan^{-1} \left(\frac{\Omega r \cos \theta}{u_o^x} \right) \quad (49)$$

From Equations 40, 41, 42, 48, and 49, the field intensity at a point on the solar equatorial plane ($\theta = 0$) is

$$B(r) = B(r_1) \left[(r_1/r)^4 + (r_1 \Omega / r u_o^x)^2 \right]^{1/2} \quad (50)$$




Figure 27. Proton Trajectory in Undisturbed Interplanetary Magnetic Field



The distance of an allowed emergence point from the line of force along which the guiding center of the proton moves is

$$\Gamma_s = \frac{m_o}{qB_s} \left(1 - \frac{u_s^2}{c^2} \right)^{-1/2} u_s \sin \beta \quad (51)$$

where B_s is the undisturbed field at the emergence point ($r = r_s$), u_s is the velocity of the scattered proton, β is the pitch angle of the trajectory, m_o is the proton rest mass, c is the velocity of light, and q is the proton charge. The field varies so slowly along the trajectory ($\lesssim 6 \times 10^{-3}$ gauss sec^{-1}) that β and u_s are very nearly adiabatic invariants of the motion. The proton arrives at the receiver from a direction at an angle β to the local field direction; the flux is axially symmetric about the field direction except for a brief initial period comparable to one gyration period. The distance (Γ_D) of the proton from its guiding center when it reaches the receiver is given by an equation similar to 51 with B_s replaced by B_R , the field at the receiver, which is found from Equations 40, 41, and 42 with $r = r_R$. The proton flux emerging from the shock at r_s with a given pitch angle $\beta < 90$ degrees is, therefore, attenuated by a factor $(r_s/r_R)^2$ at the receiver ($r = r_R$).

The time of flight of a proton from its point of emergence from the shock to the receiver is the integral of the gyration period along the trajectory,

$$t_R = \frac{\frac{2\pi}{u_s} \int_{(r_s, \phi_s)}^{(r_R, \phi_R)} \Gamma(r) \sec \beta \, d\ell}{\int_{(r_s, \phi_s)}^{(r_R, \phi_R)} d\ell} \quad (52)$$

where $d\ell$ is an element of the path (i.e., the line of force) followed by the guiding center. The line integrals in Equation 52 can be transformed to integrals over the radial coordinate r by

$$d\ell = \sec \xi \, dr = \left[\left(\frac{\Omega u_o}{r} \right)^2 + 1 \right]^{1/2} dr \quad (53)$$



The structure of the field is such that the gyroradius at any solar distance $r > r_s$ is

$$\Gamma(r) = \Gamma_s (r_s/r)^2 \left[\left(\frac{\Omega u_o x}{r} \right)^2 + 1 \right]^{1/2} \quad (54)$$

so for $r_R > r_s$


$$t_R = \frac{2m_o}{3qB_s} \left(1 - \frac{u_s^2}{c^2} \right)^{-1/2} (r_s^2 \tan \beta) \frac{\Omega^2 (u_o x)^2 \left(\frac{1}{r_s^3} - \frac{1}{r_R^3} \right) + 3 \left(\frac{1}{r_s} - \frac{1}{r_R} \right)}{(r_R^2 - r_s^2)^{1/2} - \Omega u_o x \ln \left\{ \frac{r_s [1 + (1 + r_R^2 / \Omega^2 u_o^2 x^2)^{1/2}]}{r_R [1 + (1 + r_s^2 / \Omega^2 u_o^2 x^2)^{1/2}]} \right\}} \quad (55)$$

The time of flight can be measured with reasonable accuracy from the beginning of the flash phase of the flare, when coronal heating is maximum. The results must be corrected for the time required for light emitted by the flare to reach the receiver. The time of flight t_R is much longer than the time required for light from the flare to reach the region in which acceleration occurs.

Results of this analysis are presented later in this section.

Proton Interaction with Irregularities in Medium

Magnetometers carried on interplanetary spacecraft such as Mariner 2 (Reference 56) reveal a highly disordered structure of the magnetic field in regions where the concentration of protons at energies from 10^2 to 10^4 ev is significantly greater than their concentration in the steady solar wind. These regions, referred to below as "plasma clouds," apparently result from solar activity and may be injected into space in association with the flare which produces a given energetic proton radiation event, or in association with a previous flare or eruptive prominence. The region moves outward from the sun at a velocity comparable to that of a hydromagnetic shock wave in the outer corona, i.e., about 1.0×10^6 to 1.5×10^6 m sec⁻¹, reaching 1 AU in 1.0×10^5 to 1.5×10^5 sec. If a given energetic solar proton radiation event has been preceded within one to three or four days by another such event which produced a plasma cloud, this cloud affects the



energetic proton radiation environment associated with the later event. A proton event not preceded by formation of a plasma cloud is affected by the cloud (if any) accompanying this event. The magnetic field in the plasma cloud has the effect of greatly prolonging the time required for a proton to traverse the cloud. Protons magnetically trapped in the cloud generally remain in the cloud for several hours and are transported in the cloud away from the sun.

Magnetic field measurements inside plasma clouds support the assumption that each component of the field (induction) vector \vec{B} has a Gaussian distribution about zero. The rms average scalar field intensity \bar{B} , therefore, has a χ -distribution (Reference 57) with rms average value

$$\bar{B} = \sigma_B \sqrt{3} \quad (56)$$

where σ_B is the standard deviation of a component of \vec{B} from the mean value of zero. The magnetic energy density $E_B = |\vec{B}|^2/8\pi$ has the corresponding χ^2 -distribution. A reasonable value of \bar{B} is $10 B_e$, where B_e is the scalar field intensity outside the plasma cloud; the components of B_e are given by Equations 40, 41, and 42. Plasma cloud diameters apparently range from about $0.1 R_o$ (Reference 58) to about R_o (References 59 and 60), where R_o is the heliocentric distance of the geometric centroid of the cloud.

In order to evaluate the energetic proton concentration inside a plasma cloud, the probability of reflection of an externally incident proton at the plasma boundary was calculated. The proton will be reflected if its pitch angle β about the external magnetic field (B_e) satisfies

$$\sin^2 \beta > B_e / B_i \quad (57)$$

where B_i is the field intensity in the cloud at the point of incidence, selected from the χ -distribution with rms average value \bar{B} . It can be shown (Reference 61) that the probability that (B_i/B_e) will correspond to the condition of Equation 57 is

$$P(B_i > B_e \sin^2 \beta) = 1 - \frac{B_e / \bar{B}}{2 \sqrt{6\pi} \sin^2 \beta} \sum_{n=0}^{\infty} \frac{(-3 B_e^2 / 2 \bar{B}^2 \sin^4 \beta)^n}{\left(n + \frac{3}{2}\right) \cdot n!} \quad (58)$$



The probability of absorption of the proton by the plasma,

$$P_a = 1 - P(B_i > B_e \sin^2 \beta) \quad (59)$$

is plotted in Figure 28 as a function of β for various values of (\bar{B}/B_e) in a spherical plasma cloud surrounded by a uniform field B_e .

If a proton enters the plasma, it is assumed to perform random walk on the vector space $\{\vec{r}\}$, $r < R$, where \vec{r} is the displacement of the proton in one step of the walk from its position at the beginning of the step (the position \vec{r} is measured from the center of the cloud), and R is the plasma cloud radius (Reference 62). The step length $(\delta \vec{r})$ of the random walk is variable and is selected from a χ -distribution in which the rms average scalar displacement equals the gyroradius of the proton in the magnetic field at the beginning of the step,

$$\overline{\delta r} = \frac{m_o c u}{q(c^2 - u^2)^{1/2} |\vec{u} \times \vec{B}|} \quad (60)$$

where u is the proton velocity. The field intensity B is selected from a χ -distribution as previously described; values of the angle between \vec{u} and \vec{B} are isotropically distributed. If $r > R$, the proton reaches the plasma boundary and emerges.

The distribution $r(t)$ was calculated for particles with $r < R$ after $n = 250, 500$ and 1000 steps, at proton energies of $1, 10$, and 100 Mev. The distribution of the time t_R when particles first reach $r = R$ was also calculated in order to establish the median time t_R at which one-half of the particles initially in the plasma have escaped. A practical upper bound of the effect of the plasma was sought by using $R = (2/3) R_o$, with $R_o = 1, 1.41$, and 2 AU. Figure 29 presents a typical position distribution $r(t)$; here $E = 100$ Mev, $R_o = 1$ AU, $B_e = 3\gamma$ (3×10^{-5} gauss), $\bar{B} = 30\gamma$, $n = 500$ steps, and 50 protons are assumed initially present. The sample size corresponds to a probable error of 14 percent in the concentration, which is smaller than the uncertainties in B_e and \bar{B} . The distribution of escape times t_R in this case is shown in Figure 30.

Results

The time-dependent energy distributions of protons reaching points at $1, 1.41$, and 2 AU following isotropic emergence from a hydromagnetic shock were calculated from the source function given graphically in Figure 26, using Equations 51 and 55. These results, presented as the solid curves in Figures 31, 32, and 33, are based on a hydromagnetic shock parameter




Figure 28. Probability of Absorption of Proton in Plasma




Figure 29. Space-Time Distribution of Protons in Plasma




Figure 30. Distribution of Proton Escape Times from Plasma




Figure 31. Proton Flux Rate Versus Energy at 1 AU




Figure 32. Proton Flux Rate Versus Energy at 1.41 AU




Figure 33. Proton Flux Rate Versus Energy at 2 AU



$\lambda = 4/3$ and a flare with duration 10^3 sec at the point at which the interplanetary magnetic line of force through the receiver intersects the solar chromosphere (see Figure 34). The "delay time" is the time interval following arrival of the first light from the flare at a solar distance equal to that of the receiver.

The value of λ has little effect on the proton environment (Reference 32). If the flare longitude differs by more than a few degrees from the assumed position, light from the flare cannot induce a shock in the part of the lower corona from which accelerated protons can reach the receiver. Figure 26 shows that the consequence is absence of lower-energy protons from the received spectrum. The minimum proton energy at the receiver is plotted versus solar distance in Figure 35 for various flare longitudes (at the solar equator) as observed from the receiver. The environment at energies above the cut-off is negligibly affected by the flare longitude and latitude.

The results shown in Figure 31 were integrated over time to obtain the total proton flux versus energy distribution at 1 AU, shown as the solid curve in Figure 36. The corresponding results at other solar distances may be found by dividing the result of Figure 36 by the square of the solar distance, as shown previously.

In order to include the effects of plasma clouds on the proton radiation environment, two alternative situations were considered:

1. The proton radiation event is the first to occur in at least four days. Approximately simultaneously with the flare that produces this event, a plasma cloud moves radially from the sun toward the receiver with velocity $v_c = 1160$ km/sec, the velocity of a hydromagnetic shock front with $\lambda = 4/3$ (Reference 52). The front of the cloud, with radius $R = (2/3) R_0$, reaches 1 AU in 8.5×10^4 sec, and the trailing surface passes 1 AU 1.7×10^5 sec after the flare. Protons associated with the flare are reflected or captured by the plasma near the sun; the number of protons inside the cloud decreases exponentially with median escape time \bar{t}_R , measured from the time of the flare.
2. The proton radiation event has been preceded by one that has produced a plasma cloud which, at the time of the flare causing the later event, has just reached the receiver. The protons that enter the cloud now do so at solar distances from $R_0/3$ to R_0 . The median emergence time \bar{t}_R is still measured from the time of the later flare, but now few protons emerge from the plasma before it reaches the receiver.

Results based on situations 1 and 2 are shown by dashed and dotted curves in Figure 31, 32, 33, and 36. The effect of the plasma on the total (time-integrated) radiation environment at 1 AU is not great, and is even




Figure 34. Line of Force in Quiescent Interplanetary Magnetic Field




Figure 35. Minimum Proton Energy at Receiver Versus Flare Longitude




Figure 36. Proton Flux Versus Energy at 1 AU



less at greater solar distances. The plasma does, however, considerably prolong a proton radiation event, especially at energies below 100 Mev.

Discussion of Results

The proton event model developed here exhibits several of the characteristics of observed events tabulated in Section II, including

1. The slope of the differential energy spectrum with its "knee" between 20 and 40 Mev
2. The increasing softness of the spectrum with time
3. The appearance of protons with given energy at increasing arrival angles (Reference 63)
4. The ability of flares on the western half of the visible solar hemisphere, or just east of the central meridian, or just beyond the western limb, to lead to observable proton events.

The size or total proton flux of the model event can be adjusted by altering the assumed duration of the flare.

This model is based upon the propagation of a single shock wave in the previously unshocked solar corona. After passage of the initial shock wave, the corona remains at a temperature higher than that of the unshocked corona. Secondary shocks may, therefore, produce a larger yield of energetic protons than does the initial shock, since the average proton energy before arrival of the secondary shocks is greater than in the unshocked corona. Consideration of multiple shocks may result in a greater proton flux from a flare of given duration.

This model predicts neither flares nor solar proton radiation occurrence probabilities. It is limited to a description of the proton radiation environment associated with flares bright enough and large enough to induce hydromagnetic shock waves in the corona.



V. RADIATION ENVIRONMENT EVALUATION

APPLICATION TO MARS MISSION

To illustrate the techniques discussed in previous sections, a specific mission is considered. This mission is a 1971, unmanned, one-way trip to Mars. Some of the parameters for such a mission (chosen to minimize the ΔV required) are listed in Table 12.

The solar proton event environment for this mission is chosen from the following two relationships derived earlier

$$\log_{10} \left[\int \phi_Y > 30 \text{ Mev} \right] (1971) = 8.3 + 0.0023 R_Y (1970) \quad (61)$$

$$\text{PEN} (1971) = 0.1 R_Y (1970) \quad (62)$$

where $\int \phi_Y > 30 \text{ Mev}$ is the integral particle flux (protons/cm²-year) above 30 Mev, R_Y is the annual smoothed Wolf sunspot number, and PEN is the

Table 12. Trajectory Parameters for a 1971 Mission to Mars

Date		Time After Launch (days)	Solar Distance (AU)	Longitude (degrees)
Calendar	Julian			
24 May 1971	244, 1095.5	0	1.00	242
23 June 1971	1125.5	30	1.05	273
23 July 1971	1155.5	60	1.11	301
22 August 1971	1185.5	90	1.18	326
21 September 1971	1215.5	120	1.27	346
21 October 1971	1245.5	150	1.35	6
20 November 1971	1275.5	180	1.40	22
20 December 1971	1305.5	210	1.45	38



proton event number. Equation 61 represents the major axis of the 30-Mev, 95-percent confidence ellipse (see Equation 7 and Figure 4) and, thus, corresponds to the most probable (mean) value of $\log_{10}(\int \phi_Y > 30 \text{ Mev})$.^{*} Equation 62 is the same as Equation 9; the PEN concept was illustrated in Figure 5 and Table 5.

Since R_Y has been predicted as 135 ± 54 for 1970 (Figure 2), the use of Equations 61 and 62 leads to the following values for 1971 at 1 AU:^{††}

$$\int \phi_Y > 30 \text{ Mev} = 4.1 \times 10^8 \text{ p/cm}^2 \text{ above 30 Mev}$$

$$\text{PEN} = 14$$

Based upon the observed event size distribution (Figure 23), the following combination of events (at 1 AU) appears probable:

$$1 \text{ event } 2 \times 10^8 \text{ p/cm}^2 > 30 \text{ Mev } \text{PEN} = 3$$

$$2 \text{ events } 5 \times 10^7 \text{ p/cm}^2 > 30 \text{ Mev } \text{PEN} = 2 \times 2 = 4$$

$$4 \text{ events } 2 \times 10^7 \text{ p/cm}^2 > 30 \text{ Mev } \text{PEN} = 4 \times 1 = 4$$

$$3 \text{ events } 1 \times 10^7 \text{ p/cm}^2 > 30 \text{ Mev } \text{PEN} = 3 \times 1 = 3$$

It will be noted that this combination totals to $4.1 \times 10^9 \text{ p/cm}^2 > 30 \text{ Mev}$ and a PEN of 14, as desired.

To correct for the spacecraft-sun distance, it is necessary to select probable months in which these events occur. As shown in Figure 25, July has been the most active month, with November the second most active, etc. Based upon the probability distributions of Figure 25, the following somewhat arbitrary month assignments are made for the 10 events expected in 1971:

1 event	$1 \times 10^7 \text{ p/cm}^2$	January	} at 1 AU
1 event	$5 \times 10^7 \text{ p/cm}^2$	February	
1 event	$1 \times 10^7 \text{ p/cm}^2$	March	
1 event	$2 \times 10^7 \text{ p/cm}^2$	May	
1 event	$2 \times 10^7 \text{ p/cm}^2$	July	
1 event	$2 \times 10^8 \text{ p/cm}^2$	July	
1 event	$1 \times 10^7 \text{ p/cm}^2$	August	

^{*} The curve is based upon the probability distribution of the major axis.
^{††} The curve is based upon the probability distribution of the major axis.



1 event	2×10^7 p/cm ²	September	} at 1 AU
1 event	5×10^7 p/cm ²	November	
1 event	2×10^7 p/cm ²	November	

For the hypothetical mission to Mars (Table 12) the first four events take place prior to launch and are, therefore, ignored. The expected proton fluxes for the remaining events are decreased by a factor of r_R^2 over those expected at 1 AU, where r_R is the sun-spacecraft distance in AU.

July 1971	1 event	$\int \phi = 1.6 \times 10^7$	} at spacecraft (p/cm ² >30 Mev)
July 1971	1 event	$\int \phi = 1.6 \times 10^8$	
August 1971	1 event	$\int \phi = 7.2 \times 10^6$	
September 1971	1 event	$\int \phi = 1.2 \times 10^7$	
November 1971	1 event	$\int \phi = 2.5 \times 10^7$	
November 1971	1 event	$\int \phi = 1.0 \times 10^7$	

The total expected proton flux due to solar events expected during the hypothetical mission is 2.3×10^8 p/cm² >30 Mev, as contrasted to 3.2×10^8 p/cm² >30 Mev if the entire corresponding mission had been carried out at 1 AU.

It should be pointed out that the peak flux rates are reduced by more than the factor of r_R^2 by which the integral fluxes are reduced (Figures 31 through 33). This reduction is due to an increase in the rise and decay times as r_R (the sun-spacecraft distance) increases, due to the greater path lengths from the sun and the increased dimensions of the plasma cloud. To a first approximation, the peak flux rates vary as $r_R^{-2.5}$, and so the peak flux rates expected at the spacecraft are:

July 1971	1 event	$\hat{\phi} = 200/1.28 \cong 160$	} at spacecraft (p/cm ² sec >30 Mev)
July 1971	1 event	$\hat{\phi} = 1600/1.28 \cong 1250$	
August 1971	1 event	$\hat{\phi} = 100/1.51 \cong 70$	
September 1971	1 event	$\hat{\phi} = 200/1.82 \cong 110$	
November 1971	1 event	$\hat{\phi} = 450/2.30 \cong 200$	
November 1971	1 event	$\hat{\phi} = 200/2.30 \cong 90$	



These predicted peak fluxes, when multiplied by the effective areas of the proton detectors (cm^2 - sterad), provide a basis for designing electronic circuitry to handle the expected proton detector outputs.

It will be recognized that this sort of analysis is based upon several assumptions. Among these are:

1. The proton event environment near the ecliptic plane is expected to be independent of solar longitude and latitude.
2. The monthly distribution of solar proton events is assumed to be real.
3. The integral fluxes are assumed to vary as $1/r_R^2$, where r_R is the distance from the sun.

Of these assumptions, the last is probably reasonably valid, but one of the first two is probably not. If the event environment is independent of solar latitude and longitude, there is no reason to believe that the monthly distribution is real. If, however, there is a non-uniform monthly distribution of solar proton events (at 1 AU) the solar longitude and latitude almost certainly are important. The data are not adequate to decide which, if either, of the first two assumptions is valid.

It is possible to select the numbers and sizes of solar proton events for the years 1969 through 1975 based upon the yearly smoothed Wolf sunspot number. These predictions are listed in Table 13. As before, the expected events are those that add up to the expected integral flux ($\int \phi$) above 30 Mev and the proton event number (PEN). (The events listed in Table 13 are not a unique set of events and, considering the statistical distributions about the most probable values of R_Y , PEN, and $\int \phi$, at least several hundred combinations are possible.)

The extension of this technique to a computer is straightforward. The computer would select the events from the event size distribution and the year(s) of the mission. Once the sizes of the events expected each year had been generated by a random table look-up technique, the month in which each event is expected to take place would be assigned on the basis of the monthly event distributions. This environment would then be used to calculate the mission flux expected for any preassigned spacecraft mission. By repeating this procedure several hundred times, mission flux probabilities would be generated for any desired mission. By applying this procedure to several different missions, it would be possible to calculate mission flux probabilities as a function of mission parameters.



Table 13. Solar Proton Events Expected at 1 AU for 1969-1975

Year	Ry for Previous Year	PEN	$\int \phi > 30 \text{ Mev}$ (p/cm ²)	Events Expected p/cm ² > 30 Mev
1969	128	13	3.9×10^8	1 event $\sim 2 \times 10^8$ 2 events $\sim 5 \times 10^7$ 3 events $\sim 2 \times 10^7$ 3 events $\sim 1 \times 10^7$
1970	132	13	4.0×10^8	1 event $\sim 2 \times 10^8$ 2 events $\sim 5 \times 10^7$ 4 events $\sim 2 \times 10^7$ 2 events $\sim 1 \times 10^7$
1971	135	14	4.1×10^8	1 event $\sim 2 \times 10^8$ 2 events $\sim 5 \times 10^7$ 4 events $\sim 2 \times 10^7$ 3 events $\sim 1 \times 10^7$
1972	126	13	3.9×10^8	1 event $\sim 2 \times 10^8$ 2 events $\sim 5 \times 10^7$ 3 events $\sim 2 \times 10^7$ 3 events $\sim 1 \times 10^7$
1973	108	11	3.5×10^8	1 event $\sim 2 \times 10^8$ 1 event $\sim 5 \times 10^7$ 4 events $\sim 2 \times 10^7$ 2 events $\sim 1 \times 10^7$
1974	92	9	3.2×10^8	1 event $\sim 2 \times 10^8$ 1 event $\sim 5 \times 10^7$ 3 events $\sim 2 \times 10^7$ 1 event $\sim 1 \times 10^7$
1975	77	8	3.0×10^8	1 event $\sim 2 \times 10^8$ 1 event $\sim 5 \times 10^7$ 2 event $\sim 2 \times 10^7$ 1 event $\sim 1 \times 10^7$



VI. CONCLUSIONS

The novel results of the work reported are the establishment of a solar proton radiation environmental model which includes (1) dependences on the cyclic level of solar activity and on solar distance, (2) the analytic representation of the proton flux versus energy distribution so as to include the dependence of this distribution on the total flux, and (3) the quantitative expression of the standard deviations and, hence, the probability distributions of the calculated quantities. In addition, a critical review of original research reports led to a definitive tabulation of proton radiation event and solar flare parameters. The correlation of proton radiation event and solar flare occurrence probabilities with predictable solar phenomena was established for the 1969 to 1975 period. Methods that permit calculation of not only the most probable proton radiation environment in interplanetary space flight, but also the probability of occurrence of other environmental levels have been established.

The results leave something to be desired, however. The analytical representation of the temporal dependence of the proton flux versus energy is based on data which include early and primitive measurements and which may exclude significant radiation events in years before about 1956. The association of the proton radiation environment with predictable solar phenomena is well defined but suffers from randomness of the association and of the supposedly predictable phenomena. For example, future sunspot numbers can be predicted only with some uncertainty, and the proton flux-sunspot number correlation, while significant, is not excellent.

RECOMMENDATIONS

The results are incomplete in that they portray only a portion, although perhaps the most important portion, of the complete particulate radiation environment associated with solar flares. Additional work is recommended to provide a fuller description of this environment. The recommended work primarily concerns inclusion of particle types other than protons, evaluation of directional distributions of the radiation flux, extension of the solar distance range, and improvement of the method of evaluating environmental levels on interplanetary missions.

The charged particulate radiation emitted in association with solar flares may contain, in addition to protons, electrons, alpha particles, and heavy nuclei (References 64, 65, and 66). It has been shown (References 64



and 67) that alpha particle radiation may produce greater material and biological damage than protons produce. Electrons may also be a significant part of the radiation environment of exposed or thinly protected spacecraft components. Radiation instrumentation intended to accomplish comprehensive studies of space radiation and solar activity should be designed in light of the most probable levels of all types of particles composing the radiation environment; therefore, the electron and alpha particle fluxes versus energy, time after a flare, solar activity level, and solar distance should be evaluated. If feasible, similar results should be obtained for the flux of heavy nuclei (lithium to iron). Results of experimental measurements of electrons and alpha particles in space should be supplemented by analyses based on particle acceleration by hydromagnetic shocks, synchrotron radiation energy loss of electrons in the solar magnetic field, and diffusion of energetic particles in local clouds of disordered plasma in the interplanetary medium.

The analyses of hydromagnetic shock acceleration and proton transport presented in Section IV can be used to provide values of the distribution of solar proton arrival angles versus time, proton energy, and solar direction (Reference 54). These results agree well with observed angular distributions in the vicinity of the earth (Reference 68) and also with earlier models of the anisotropy of solar proton radiation (Reference 69). The directional distribution of the radiation is significant in spacecraft shielding analysis and in the design of particle spectrometer arrays which measure the angular distribution of the charged particle flux. Therefore, the solar radiation directional distribution should be evaluated for protons, alpha particles, and electrons as a function of time after a flare, stage of the sunspot number cycle, particle energy, and solar distance. This work will reveal the dependence of the degree of anisotropy on solar distance.

The solar distance interval from 1 to 2 AU contains most unmanned vehicle trajectories to Mars. Manned missions to Mars reach aphelion distances beyond 2 AU before returning to earth; any flights to the major asteroids or Jupiter extend to as much as 5 AU. Approaching the sun, Venus missions which have immediate interest reach perihelion distances near 0.7 AU, while future solar probes and Mercury missions may reach solar distances as small as 0.3 AU. It is, therefore, worthwhile to extend the proton radiation models developed here, as well as the recommended studies of electron and alpha particle environments and of anisotropy, to the solar distance range from 0.3 to 5 AU. The analyses below 1 AU will be very straightforward because of the diminishing importance of irregularities in the interplanetary medium, while results beyond 2 AU will admittedly suffer in reliability because of the poorly known extent of solar magnetic fields and the solar wind into the interstellar medium. The importance of the galactic particle radiation environment also increases with increasing solar distance, both absolutely and with respect to the solar radiation.



The application of the present and recommended analyses to the evaluation of the radiation environment on space missions can be improved by automation. It is recommended that a computer program be prepared which uses Monte Carlo techniques to select randomly from probability distribution tables the dates and particle fluxes of the solar particulate radiation events encountered on a specified trajectory. If the same mission is "flown" repeatedly, the probability distribution of radiation environment profiles can be established.

The work presented here provides a valuable technique for estimating the expected solar proton radiation environment in interplanetary space. If this work is supplemented by the recommended effort, the planning of space missions and the design of space vehicles will benefit from a quantitative statistical description of present knowledge of all significant aspects of the radiation environment associated with solar activity.



VII. REFERENCES

(Additional bibliographical listings are given in Appendix A; references listed here are those cited in the text).

1. Models for Space Environmental Hazards: Radiation, NASA Manned Spacecraft Center unpublished report (31 January 1963).
2. Bailey, D.K. Time Variations of the Energy Spectrum of Solar Cosmic Rays in Relation to the Radiation Hazard in Space, J. Geophys. Res. 67, 391 (1962).
3. Modisette, J.L., T.M. Vinson, and A.C. Hardy. Model Solar Proton Environments for Manned Spacecraft Design, NASA Technical Note D-2746 (April 1965).
4. Webber, W.R. An Evaluation of the Radiation Hazard due to Solar Particle Events, Report D2-90469, The Boeing Company (1963).
5. McDonald, F.B. Solar Proton Manual, NASA Technical Report TR-R-169 (1963).
6. Lewis, L.R. et al. Solar Flare Radiation Survey, Report RTD-TDR-63-3044, U.S. Air Force Weapons Laboratory (1963).
7. Hess, W.N. et al. AAS-NASA Symposium on the Physics of Solar Flares, NASA Report SP-50 (1964).
8. Malitson, H.H. Table of Solar Proton Events, Appendix of Solar Proton Manual, F. B. McDonald, ed, NASA TR-R-169 (1963).
9. Malitson, H.H. and W. R. Webber. A Summary of Solar Cosmic Ray Events, Chapter 1 of Solar Proton Manual, F. B. McDonald, ed, NASA TR-R-169 (1963).
10. Fletcher, J. D. Solar Radio Emission as a Criterion for Solar Proton Event Warning, AIAA Jour. 2, 2193 (1964).
11. Pomerantz, M.A. Cosmic Ray Observations at Very High Altitudes During Periods of Intense Solar Activity, Phys. Rev. 102, 870 (1956).



12. Goedeke, A. D. and A. J. Masley. Observations in the Antarctic of Solar Cosmic Ray Events in 1962 and 1963, J. Geophys. Res. 69, 4166 (1964).
13. Webber, W.R. A Review of Solar Cosmic Ray Events, AAS-NASA Symposium on the Physics of Solar Flares, NASA SP-50 (1963) p. 215.
14. Fitchel, C.E., D.E. Guss, and K.W. Ogilvie. Details of Individual Solar Particle Events, Solar Proton Manual, F.B. McDonald, ed., NASA TR-R-169 (1963) p. 19.
15. Symposium on the July 1959 Events and Associated Phenomena, International Union of Geodesy & Geophysics, Helsinki, Finland, Monograph No. 7 (July 1960).
16. Webber, W.R. and H.H. Malitson. A Summary of Solar Cosmic Ray Events, Solar Proton Manual, Report X-611-62-122, NASA Goddard Space Flight Center.
17. Bailey, D.K. Catalog of Polar Cap Absorption Events (1952 to 1961) From VHF Ionospheric Scatter Observations, with Details of Possible Related Preceding Solar Flares and Short Wave Radio Fadeouts, NBS Report, Boulder, Colo.
18. Foelsche, T. Estimates of Radiation Doses in Space on the Basis of Current Data, Life Sciences and Space Research, a session of the Third International Space Science Symposium, Washington D.C., April 30-May 9, 1962 (1963) p. 48.
19. Yoshida, S. et al. Propagation of Solar Particles Deduced From Measurements with Explorer VII, Space Research III, Proceedings of the Third International Space Science Symposium, Washington D.C., May 2-8, 1962 (1963) p. 608.
20. Bailey, D.K. Polar Cap Absorptions, Planet. Space Science 12, 495 (1964).
21. Wasson, J. T. Radioactivity Produced in Discoverer XVII by November 12, 1960, Solar Flare Protons, J. Geophys. Res. 66, 2659 (1961).
22. Biswas, S., C.E. Fitchel, and D.E. Guss. Study of H, He, and Heavy Nuclei in November 12, 1960 Solar Cosmic Ray Event, Phys. Rev. 128, 2756 (1962).




23. Ogilvie, K. W., D. A. Bryant, and L. R. Davis. Rocket Observations of Solar Protons during the November 1960 Events, 1, J. Geophys. Res. 67, 929 (1962).
24. Ney, E. P. and W. A. Stein. Solar Protons, Alpha Particles, and Heavy Nuclei in November 1960, J. Geophys. Res 67, 2087 (1962).
25. de Feiter, L. D., A. Freon, and J. P. LeGrand. The Cosmic Ray Flares of November 12 and 15, 1960, Space Res. 2, 776 (1961).
26. Kodama, M. and M. Kitamura. Some Features of the November 1960 Events as Inferred from Cosmic Rays, Space Res. 2, 766 (1961).
27. Pomerantz, M. A., S. P. Duggal, and K. Nagashima. The Unusual Cosmic Ray Intensity Increases on November 12, 1960, Space Res. 2, 788 (1961).
28. Conforto, A. M., and N. Iucci. Cosmic Ray Events in November 1960, Space Res. 2, 787 (1961).
29. Lockwood, J. A. and M. A. Shea. Variations of the Cosmic Radiation in November 1960, J. Geophys. Res. 66, 3083 (1961).
30. Roederer, J. G. et al. Cosmic Ray Phenomena During the November 1960 Solar Disturbances, J. Geophys. Res. 66, 1603 (1961).
31. Ogilvie, K. W. and D. A. Bryant. Solar Proton Spectra in the events of November 12 and 15, 1960, J. Geophys. Res. 69, 393 (1964).
32. Masley, A. J. and A. D. Goedeke. Complete Dose Analysis of the November 12, 1960 Solar Cosmic Ray Event, Life Sciences and Space Research, a session of the Third International Space Science Symposium, R. B. Livingston, A. A. Imshenetsky & G. A. Derbyshire, ed., North Holland (1963).
33. Krimigis, S. M., J. A. Van Allen, and L. A. Frank. Observations with University of Iowa Equipment on Mariner IV, May-October 1965, Report 65-36, University of Iowa (1965).
34. Van Allen, J. A. Solar Cosmic Ray Events, January-September 1963, Trans. Am. Geophys. Union 45, 351 (1964).
35. Cline, T. C. et al. Explorer 14 Observations of Solar Proton Events and Interplanetary Particle Events in 1962 and 1963, Trans. Am. Geophys. Union 46, 129 (1965).



36. Weddell, J. B. Prediction of Probability of Occurrence of Solar Flares, Astronautica Acta 10, 340 (1964).
37. Anderson, K. A. Preliminary Study of Prediction Aspects of Solar Cosmic Ray Events, NASA Technical Note D-700 (1961).
38. Waldmeier, M. Ergebnisse und Probleme der Sonnenforschung, Zer Aufl., Geest and Portig, Leipzig (1955), pp. 139-157.
39. Churchill, R. V. Fourier Series and Boundary Value Problems, McGraw-Hill Book Company, New York (1941), pp. 88-91.
40. T. K. Oliver, private communication.
41. Minnis, C. M. An Estimate of the Peak Sunspot Number in 1968, Nature 186, 462 (1960).
42. King-Hele, D. C. Prediction of the Dates and Intensities of the Next Two Sunspot Maxima, Royal Aircraft Establishment report, Farnborough, England (November 1965).
43. Aitken, A. C. Statistical Mathematics, Oliver and Boyd Ltd., Edinburgh (1949), p. 88.
44. Callender, R. H., J. R. Manzano, and J. R. Winckler. The Response of High Altitude Ionization Chambers During the 1954 - 1965 Solar Cycle, J. Geophys. Res. 70, 3189 (1965).
45. Compilations of Solar-Geophysical Data, Central Radio Propagation Laboratory Bulletin, Series F, Part B, National Bureau of Standards, Boulder, Colorado (1956-1965).
46. Severnyi, A. B. Some Results of Investigations of Nonstationary Processes on the Sun, Soviet Astronomy - AJ 1, 668 (1957).
47. Howard, R. Observations of the Solar Magnetic Field, Astrophys. J. 130, 193 (1959).
48. Legrand, J. -P. Possibilités de Prévision des Phénomènes Solaires et de Leurs Répercussions sur le Globe par l' Étude de l' Intensité du Rayonnement Cosmique, Comptes Rend. 248, 70 (1958).
49. Hodder, D. T. NAA S&ID report (unpublished).
50. Haffner, J. W. Shielding Analysis of 1956-1961 Solar Proton Event Data, NAA S&ID, SID 64-1295 (1964).



51. Haffner, J. W. An Attenuation Kernel for the Bailey Model Event, Trans. Am. Nucl. Soc. 7, 16 (1964).
52. Parker, E. N. Interplanetary Dynamical Processes, Interscience Publishers, New York (1963), pp. 73-80, 99-105.
53. Weddell, J. B. Interaction of Hydromagnetic Shocks with Charged Particles in the Solar Corona, NAA S&ID, SID 64-1500 (August 1964).
54. Weddell, J. B. Solar Proton Event Production by Hydromagnetic Shocks, Planet. Space Sci. 13, 789 (1965).
55. Parker, E. N. Dynamics of the Interplanetary Gas and Magnetic Fields, Astrophys. J. 128, 664 (1958).
56. Coleman, P. J. et al. Interplanetary Magnetic Fields, Science 138, 1099 (1962).
57. Menzel, D. H. Fundamental Formulas of Physics, Dover Publications, Inc., New York (1960), Vol. 1, p. 110.
58. Neugebauer, M. and C. W. Snyder. Solar Plasma Experiment, Science 138, 1095 (1962).
59. Dvoryashin, A. S. Proton Flares from 1957 to 1961 and the Geometry of Interplanetary Magnetic Field, NASA TT F-226 (November 1964); Izv. Krymskoy Astrof. Obs. 28, 293 (1962).
60. Wilcox, J. M. and N. F. Ness. Quasi-Stationary Corotating Structure in the Interplanetary Medium, J. Geophys. Res. 70, 5793 (1965).
61. Weddell, J. B., and D. A. McQuarrie. Interaction of Energetic Solar Protons with Disordered Magnetic Fields, Trans. Am. Geophys. Un. 47, 152 (1966).
62. Bharucha-Reid, A. T. Elements of the Theory of Markov Processes and Their Applications, McGraw-Hill Book Company, Inc., New York (1960), pp. 45-46.
63. McCracken, K. G. The Cosmic Ray Flare Effect. 2. The Flare Effects of May 4, November 12, and November 15, 1960, J. Geophys. Res. 67, 434 (1962).
64. Sheldon, W. R., and S. B. Curtis. Radiation Hazard from Heavy Particles in Solar-Flare Events, Bull. Am. Phys. Soc., Series II, 9, 141 (1964).



65. Van Allen, J. A., and S. M. Krimigis. Impulsive Emission of ~40-kev Electrons from the Sun, J. Geophys. Res. 70, 5737 (1965).
66. Arnoldy, R. L., S. R. Kane, and J. R. Winckler. Observations of Energetic Electrons with the Ogo A Ionization Chamber from the June 5, 1965, Solar Flare, Trans. Am. Geophys. Union 47, 154 (1966).
67. Haffner, J. W. The Role of Alpha Particles in Shielding Against Solar Event Radiation, NAA S&ID, SID 64-1297 (1964).
68. McCracken, K. G. The Cosmic Ray Flare Effect. 2. The Flare Effects of 4 May, 12 November, and 15 November 1960, J. Geophys. Res. 67, 435 (1962).
69. Stazer, A. K., A Model Anisotropic Solar Proton Event, presented at American Institute of Aeronautics and Astronautics, Aerospace Sciences meeting, New York (22-24 January 1964).



APPENDIX A. BIBLIOGRAPHY

This bibliography cites all significant documents located in the literature survey described in Section II of this report. The first part of the bibliography lists references to specific solar flares and flare-associated proton radiation events and magnetic disturbances; this list is in chronological order of the natural events described. Each of the following parts is topical, in chronological order of publication. The topics are prediction of solar flares, origin of solar flares, solar radio bursts, source of solar flare particles, propagation of solar flare particles, cosmic ray (i.e., galactic particle) effects, energy spectra of solar flare particles, polar cap absorptions, geomagnetic disturbances related to solar flares, ionization and reactions in the earth's atmosphere, and miscellaneous references.

SPECIFIC SOLAR FLARES AND PROTON RADIATION EVENTS

November 1949

Rose, D.C. On the Sudden Increase in Cosmic Ray Intensity on November 19, 1949, Phys. Rev. 78, 181 (1950).

Forbush, S.E., T.B. Stinchomb, and M. Schein. On the Extraordinary Increase of Cosmic Ray Intensity on November 19, 1949, Phys. Rev. 78, 323 (1950).

Forbush, S.E., T.B. Stinchomb, and M. Schein. The Extraordinary Increase of Cosmic Ray Intensity on November 19, 1949, Phys. Rev. 79, 501 (1950).

Clay, J., and H.F. Jorgen. The Solar Flare of November 19, 1949 and Cosmic Rays, Phys. Rev. 79, 908 (1950).

August 1950

Naugle, J.E., and G.W. Anderson. Rates of Star Formation and the Flux of Primary Heavy Nuclei During the Solar Flare of August 2, 1950, Phys. Rev. 81, 314 (1951).

Dodson, H.W., and R.W. Donselman. The Eruptive Prominence of August 7, 1950, Astrophys. J. 113, 519 (1951).

May 1951

Bartlett, T.J., B. Witte, and W.O. Roberts. Eruptive Prominence Associated with Limb Flare of May 8, 1951, Astrophys. J. 117, 292 (1953).

August 1954

Dodson, H.W. Development of an Active Solar Region, August 20-26, 1954, Astron. J. 61, 2 (1956).

February 1956

Dorman, L.I. et al. Observation of the Large Cosmic Ray Increase of February 23, 1956 in the USSR, Nuclear Phys. 1, 585 (1956).

Bachelet, F. and A.M. Conforto. Increase of Cosmic Ray Intensity Associated with the Solar Flare of February 23, 1956, Nuovo Cimento, Series X, 3, 1153 (1956).

Filosofo, I. et al. The Increase in the Total Cosmic Ray Intensity and in the Positive Excess due to the Solar Flare of 23rd February 1956, Nuovo Cimento, Series X, 3, 1112 (1956).

Belrose, J.S., M.H. Davenport, and K. Weekes. Some Unusual Radio Observations Made on 23 February 1956, J. Atmos. Terres. Phys. 8, 281 (1956).

Cranshaw, T.E., W. Galbraith, and N.A. Porter. Cosmic Ray Observations During the Flare of 23 February 1956, J. Atmos. Terres. Phys. 8, 274 (1956).

Ellison, M.A. and J.H. Reid. A Long-Wave Anomaly Associated with the Arrival of Cosmic Ray Particles of Solar Origin on 23 February 1956, J. Atmos. Terres. Phys. 8, 291 (1956).

Gold, T. and D.R. Palmer. The Solar Outburst, 23 February 1956 - Observations of the Royal Greenwich Observatory, J. Atmos. Terres. Phys. 8, 287 (1956).

Lambie, M. and H. Elliot. Cosmic Rays and the Solar Flare of 23 February 1956, J. Atmos. Terres. Phys. 8, 277 (1956).

Marsden, P.L. et al. Variation of Cosmic Ray Nuclear Intensity During the Disturbance of 23 February 1956, J. Atmos. Terres. Phys. 8, 278 (1956).

Ehmert, A. and G. Pfozter. A New Outburst of Solar Cosmic Rays, Z. Naturforsch. 11a, 322 (1956).



Sittkus, A. The Anomalous Increase of the Cosmic Radiation on 23 February 1956, Z. Naturforsch. 11a, 604 (1956).

Menzel, D.H. et al. The Record Prominence of 10 February 1956, Astron. J. 61, 186 (1956).

Brode, R.B. and A. Goodwin. Extraordinary Increase of Cosmic Radiation on February 23, 1956, Phys. Rev. 103, 377 (1956).

Lockwood, J.A. et al. Cosmic Ray Neutron Intensity Increase Associated with Solar Flare of February 23, 1956, Phys. Rev. 103, 247 (1956).

Meyer, P., E.N. Parker, and J.A. Simpson. Solar Cosmic Rays of February 1956 and Their Propagation Through Interplanetary Space, Phys. Rev. 104, 768 (1956).

Winckler, J.R. Cosmic Ray Increase at High Altitude on February 23, 1956, Phys. Rev. 104, 220 (1956).

Dorman, L.I. et al. A Large Outburst of Cosmic Ray Intensity on February 23, 1956 Observed in Moscow, Sverdlovsk, Tbilisi, and Cape Schmidt, Soviet Physics - Doklady 108, 809 (1956).

Lied, F. Ionospheric Absorption Observed on the 23rd February 1956 at Kjeller and Tromsu, J. Atmos. Terres. Phys. 10, 48 (1957).


Van Allen, J.A. and J.R. Winckler. Spectrum of Low-Rigidity Cosmic Rays During Solar Flare of February 23, 1956, Phys. Rev. 106, 1072 (1957).

Eckhartt, O. The Electromagnetic State in Interplanetary Space During the Flare of February 23, 1956, Nuovo Cimento, Series X, Supplement 8, 202 (1958).

Pfotzer, G. Latitude Effect of Solar Cosmic Radiation on February 23, 1956 with Respect to the Problem of Effective Geomagnetic Coordinate System, Nuovo Cimento, Series X, Supplement 8, 220 (1958).

Pfotzer, G. On the Separation of Direct and Indirect Fractions of Solar Cosmic Radiation on February 23, 1956 and on the Difference in Steepness of Momentum Spectrum of These Two Components, Nuovo Cimento, Series X, Supplement 8, 180 (1958).

Goldberg, L., O.C. Mohler, and E.A. Muller. The Profile of H α During the Limb Flare of February 10, 1956, Astrophys. J. 127, 302 (1958).

June 1956

Zirin, H. On the Enhancement of Certain Helium Lines in the Limb Flare of June 24, 1956, Astrophys. J. 126, 159 (1957).

August 1956

McCracken, K.G. Production of Cosmic Radiation by a Solar Flare on August 31, 1956, Nuovo Cimento, Series X, 13, 1074 (1959).

Kriusky, L. On the Flare of August 31, 1956, Connected with the Ejection of Cosmic Ray After Y-Shaped Phase, Astron. Inst. Czecho. Bull. 15, 75, (1964).

November 1956

Tandberg-Hanssen, E. and H. Zirin. Flares and Active Prominences I - The Loop Prominences of November 12 and 22, 1956, Astrophys. J. 129, 408 (1959).

December 1956

Zirin, H. Flares and Active Prominences II - A Remarkable Limb Flare, December 18, 1956, Astrophys. J. 129, 414 (1959).

Tandberg-Hanssen, E. Flares and Active Prominences III - The Difference Between the Surge and Loop Prominences of December 19, 1956, Astrophys. J. 130, 202 (1959).

August 1957

Anderson, K.A. Soft Radiation Events at High Altitude During the Magnetic Storm of August 29-30, 1957, Phys. Rev. 111, 1397 (1958).

McNarry, L.R. The Observation of a Solar Event in White Light from Resolute N.W.T. on August 30, 1957, J. Roy. Astron. Soc. Can. 54, 273 (1960).

Anderson, K.A. Solar Cosmic Ray Events During Late August 1957, J. Geophys. Res. 69, 1743 (1964).

September 1957

Jefferies, J. T., E. v. P. Smith, and H. J. Smith. The Flare of September 18, 1957, Astrophys. J. 129, 238 (1959).

October 1957

McCracken, K.G. and N.R. Parsons. Unusual Cosmic Ray Intensity Fluctuations Observed at Southern Stations During October 21-24, 1957, Phys. Rev. 112, 1798 (1958).

February 1958

Axford, W.I. and G.C. Reid. Polar Cap Absorption and the Magnetic Storm of February 11, 1958, J. Geophys. Res. 67, 1692 (1962).

Dorman, L.I. and O.I. Inozertseva. Disturbed Solar Diurnal Variation and Anomalous Increase in the Intensity of Cosmic Rays During the Magnetic Storm of February 11, 1958, in Section VII of Cosmic Rays (Results of Researches on the Program of the International Geophysical Year), S.N. Vervov and L.I. Dorman, eds., USSR Academy of Sciences (1963), p. 62.

Hakura, Y. and M. Nagai. Synthetic Study of Severe Solar Terrestrial Disturbances on February 9-12, 1958, Radio Res. Lab. J. 11, 197 (1964).

Ganziliz, C. and L.R. Terrazas y Gradiela, The Solar Flare of Feb. 9, 1958, Observatorias Tanantzentla y Zacuboya, Boletin 3, 325 (1964). (In Spanish)

March 1958

Freier, P.S., E.P. Ney, and J.R. Winckler. Balloon Observation of Solar Cosmic Rays on March 26, 1958, J. Geophys. Res. 64, 685 (1959).

Rourke, G.F. Minor PCA Events During March 1958, J. Geophys. Res. 66, 4316 (1961).

July 1958

Rymko, N.P., V.F. Tulinov, and A.N. Charakhchyan. A Case of a Sharp Increase in Cosmic Ray Intensity in the Stratosphere, Soviet Physics-JETP 9, 1202 (1959).

Hultqvist, B., J. Aarons, and J. Ortner. Effects of the Solar Flares of 7 July 1958, Tellus 11, 319 (1959).

Letfus, V. Preliminary Curves of Growth for Iron Emission Lines in the Flare of July 20, 1958, Astron. Inst. Czech. Bull. 15, 211 (1964).

August 1958

Anderson, K. A. et al. Observations of Low-Energy Solar Cosmic Rays from the Flare of 22 August 1958, J. Geophys. Res. 64 1133 (1959).

Odintsova, I. N. Estimation of the Spectrum of the Solar Ionizing Radiation During the Flare of August 22, 1958 Using Data from the Vertical Probing of the Ionosphere; Geomagnetizm i Aeronomiia 4, 1035 (1964).

September 1958

Ondoh, T. Variations of the Earth's Exosphere and Geomagnetic Disturbances During the Geomagnetic Storm of 4 September 1958, J. Atmos. Terres. Phys. 22, 233 (1961).

May 1959

Bachelet, F. et al. The Cosmic Ray Storm of May 11, 1959, Nuovo Cimento, Series X, 13, 1055 (1959).

Eriksen, K. W., O. Holt, and B. Landmark. A Note on the Polar Absorption Event of 11-18 May 1959, J. Atmos. Terres. Phys. 18, 78 (1960).

Manzano, J. R., J. G. Roederer, and O. R. Santochi. Cosmic Ray Intensity Variations During the Magnetic Storm in May 1959, Nuovo Cimento, Series X, 18, 136 (1960).

Winckler, J. R. and P. D. Bhavsar. Low-Energy Solar Cosmic Rays and the Geomagnetic Storm of May 12, 1959, J. Geophys. Res. 65, 2637 (1960).

Biswas, S. Forbush Decreases of the Flux of Heavy Primary Nuclei of Cosmic Rays on May 12 and July 12, 1959, J. Geophys. Res. 66, 2653 (1961).

June 1959

Rymko, N. P., V. F. Tulinov, and A. N. Charakhchyan. A Case of a Sharp Increase in Cosmic-Ray Intensity in the Stratosphere, Soviet Physics--JETP 9, 1202 (1959).

July 1959

Brown, R. R. and R. G. D'arcy. Observations of Solar Flare Radiation at High Latitude During the Period July 10-17, 1959, Phys. Rev. Ltrs. 3, 390 (1959)



Carmichael, H. and J. F. Steljis. Unusual Cosmic Ray Fluctuations on July 17 and 18, 1959, Phys. Rev. Ltrs. 3, 392 (1959).

Leinbach, H. and G. C. Reid. Polar Cap Absorption During the Solar Cosmic Ray Outbursts of July 1959, International Union of Geodesy and Geophysics, Symposium on the July 1959 Events and Associated Phenomena, Helsinki, July 1960; Paris: Institut Geographique National (1960), pp. 145-150.

Cattani, D. and M. Galli. Cosmic Ray Storms of July 1959, Nuovo Cimento, Series X, 16, 765 (1960).

Roederer, J. G. et al. Preliminary Report on Cosmic Ray Intensity During Magnetic Storms in July 1959, Nuovo Cimento, Series X, 18, 120 (1960).

Bailey, D. K. and M. A. Pomerantz. The Cosmic-Ray Increase of 17 July, 1959, Can. J. Phys. 38, 333 (1960).

Wilson, B. G., D. C. Rose, and M. A. Pomerantz. The Unusual Cosmic Ray Events of July 17-18, 1959, Can. J. Phys. 38, 328 (1960).

Dodson, H. W. and E. R. Hedeman. Flares of July 16, 1959, Astron. J. 65, 51 (1960).

Ehmert, A. et al. Observations of Solar Flare Radiation and Modulation Effects at Balloon Altitudes, July 1959, J. Geophys. Res. 65, 2685 (1960).

McCracken, K. G. and R. A. R. Palmeira. Comparison of Solar Cosmic Ray Injections Including July 17, 1959 and May 4, 1960, J. Geophys. Res. 65, 2673 (1960).

Howard, R. and H. W. Babcock. Magnetic Fields Associated with the Solar Flare of July 16, 1959, Astrophys. J. 132, 218 (1960).



Athay, R.G. The Cosmic Ray Flares of July 1959 and November 1960 and Some Comments on Physical Properties and Characteristics of Flares, Space Res. 2, 837 (1961).

Hakura, Y. and T. Goh. Unusual Solar-Terrestrial Events in July 1959, Space Res. 2, 803 (1961).

Earl, J.A. Balloon Measurement of Solar Cosmic Rays at Fort Churchill, Canada, During July 1959, J. Geophys. Res. 66, 3095 (1961).

Winckler, J.R., P.D. Bhavsar, and L. Peterson. The Time Variations of Solar Cosmic Rays During July 1959 at Minneapolis, J. Geophys. Res. 66, 995 (1961).

Ghielmetti, H.S. The Spectrum and Propagation of Relativistic Solar Flare Particles During July 17-18, 1959, J. Geophys. Res. 66, 1611 (1961).

Biswas, S. The Flux of Heavy Nuclei in the July 10, 1959 Flare, J. Geophys. Res. 67, 2613 (1962).

Freier, P.S. The Measurement of the Cutoff Rigidity at Minneapolis Using Solar Protons and Particles from the July 14, 1959 Flare, J. Geophys. Res. 67, 2617 (1962).

October 1959

Krivsky, L. et al. Solar Flare Connected with an Increase of Intensity of Cosmic Rays, Nuovo Cimento, Series X, 15, 695 (1960).

November 1959

Eriksen, K.W., O. Holt, and B. Landmark. A Note on the Polar Absorption Event of 11-18, 1959, J. Atmos. Terres. Phys. 18, 78 (1960).

April 1960

Chinburg, D. L. Great Magnetic Storm of March 31-April 3, 1960, J. Geophys. Res. 65, 2206 (1960).

Van Allen, J. A. and W. C. Lin. Outer Radiation Belt and Solar Proton Observations with Explorer VII During March-April 1960, J. Geophys. Res. 65, 2996 (1960).

Masley, A. J., T. C. May, and J. R. Winckler. Analysis of Balloon Observations during the April 1960 Solar Cosmic Ray Events, J. Geophys. Res. 67, 3243 (1962).

May 1960

Hladky, J. et al. Cosmic Ray Event of May 4, 1960, Nuovo Cimento, Series X, 17, 610 (1960).

Marsden, P. L., J. B. Crowden, and C. J. Hatton. Increase in the Intensity of Cosmic Radiation on May 4, 1960, Nuovo Cimento, Series X, 17, 977 (1960).

Santochi, O. R., J. R. Manzano, and J. G. Roederer. Cosmic Ray Increase on May 4, 1960, Nuovo Cimento, Series X, 17, 119 (1960).

Rose, D. C. The Sudden Increase in Cosmic Ray Intensity of May 4, 1960, Can. J. Phys. 38, 1224 (1960).

Anderson, J. C. et al. The Solar Cosmic-Ray Outburst of May 4, 1960, J. Geophys. Res., 65, 3889 (1960).

Brode, R. B., R. R. Brown, and W. R. Steiger. Solar Flare Cosmic-Ray Increase of May 4, 1960, J. Geophys. Res. 65, 4200 (1960).

McCracken, K. G., and R. A. R. Palmeira. Comparison of Solar Cosmic Ray Injections Including July 17, 1959 and May 4, 1960, J. Geophys. Res. 65, 2673 (1960).

Trainor, J. H., M. A. Shea, and J. A. Lockwood. Increase of the Nucleonic Intensity on May 4, 1960, J. Geophys. Res. 65, 3001 (1960).

Standil, S., R. P. Bukata, and F. K. Chin. The Sudden Increase in the Photon Component of Cosmic Radiation on May 4, 1960, Can. J. Phys. 39, 229 (1961).



Biswas, S. and P. S. Freier. The High-Energy Cosmic-Ray Flare of May 4, 1960. 2. Emulsion Measurements, J. Geophys. Res. 66, 1029 (1961).

Maeda, K., V. L. Patel, and S. F. Singer. Solar Flare Cosmic-Ray Event of May 4, 1960, J. Geophys. Res. 66, 1569 (1961).

Winckler, J. R., A. J. Masley, and T. C. May. The High-Energy Cosmic-Ray Flare of May 4, 1960. 1. High-Altitude Ionization and Counter Measurements, J. Geophys. Res. 66, 1023 (1961).

McCracken, K. G. The Cosmic-Ray Flare Effect, 2. The Flare Effects of May 4, November 12, and November 15, 1960, J. Geophys. Res. 67, 435 (1962).

Chasson, R. L. Families of Flare Particles on May 4, 1960, Proceedings of 8th International Conference on Cosmic Rays, Vol. 1 (Solar Particles and Sun-Earth Relations), Tata Institute of Fundamental Research, Bombay (1964), pp. 161-165.

July 1960

Wilson, B. G., D. C. Rose, and M. D. Wilson. Small Cosmic Ray Increases Measured at Ground Level September 3, 1960, July 18, and July 20, 1960, Can. J. Phys. 40, 540 (1962).

August 1960

Freier, P. S., and C. J. Waddington. Intensity of 80 to 200 Mev Protons over Fort Churchill on August 26, 1960, J. Geophys. Res. 70, 2111 (1965).

September 1960

Davis, L. R. et al. Rocket Observations of Solar Protons on September 3, 1960, Phys. Rev. Ltrs. 6, 492 (1961).

Winckler, J. R. et al. Delayed Propagation of Solar Cosmic Rays on September 3, 1960, Phys. Rev. Ltrs. 6, 488 (1961).

Smith, E. P. and P. S. McIntosh. Spectrum of the Flare of September 2, 1960, Astron. J. 66, 54 (1961).



Wilson, B. G., D. C. Rose, and M. D. Wilson. Small Cosmic-Ray Increases Measured at Ground Level, September 3, 1960, July 18, and July 20, 1960, Can. J. Phys. 40, 540 (1962).

Biswas, S., P. S. Freier, and W. Stein. Solar Protons and Alpha Particles from the September 3, 1960 Flares, J. Geophys. Res. 67, 13 (1962).

Earl, J. A. Cloud-Chamber Observations of Solar Cosmic Rays over Minneapolis on September 4, 1960, J. Geophys. Res. 67, 2107 (1962).

Fitchel, C. E., D. A. Kniffen, and K. W. Ogilvie. September 26, 1960 Solar Cosmic Ray Event, J. Geophys. Res. 67, 3669 (1962).

Winckler, J. R. and P. D. Bhavsar. The Time Variations of Solar Cosmic Rays during the September 3, 1960 Event, J. Geophys. Res. 68, 2099 (1963).

Smith, E. P. Spectrum of the Flare of September 2, 1960, Astrophys. J. 137, 580 (1963).

November 1960

Ogilvie, K. W., D. A. Bryant, and L. R. Davis. Rocket Observations of Solar Protons during the November 1960 Event, Proceedings International Conference on Cosmic Rays and the Earth Storm, Kyoto, September 1961. II. Joint Sessions. Physical Society of Japan, Tokyo (1962), pp. 317-319.

Conforto, A. M., and N. Iucci. Cosmic Ray Events in November 1960, Space Res. 2, 787 (1961).

de Feiter, L. D., A. Freon, and J. P. LeGrand. The Cosmic Ray Flares of November 12 and 15, 1960, Space Res. 2, 776 (1961).

Ehmert, A., H. Erbe, and G. Pfozter. Peculiarities of the Outburst of Solar High Energy Particles on November 1960, Space Res. 2, 778 (1961).

Kodama, M., and M. Kitamura. Some Features of the November 1960 Events as Inferred from Cosmic Rays, Space Res. 2, 766 (1961).

Ortner, J., A. Egeland, and B. Hultquist. The Great Earth Storms of November 1960 as Observed at Kiruna Geophysical Observatory, Space Res. 2, 722 (1961).

Athay, R. G. The Cosmic Ray Flares of July 1959 and November 1960 and Some Comments on Physical Properties and Characteristics of Flares, Space Res. 2, 837 (1961).



Pomerantz, M. A., S. P. Duggal, and K. Nagashima. The Unusual Cosmic Ray Intensity Increases on November 12, 1960, Space Res. 2, 788 (1961).

Hanson, R. T. Cosmic Ray Flare of November 20, 1960, Phys. Rev. Ltrs. 6, 260 (1961).

Manzano, J. R. et al. Cosmic Ray Phenomena during the November 1960 Solar Disturbances, Notas de Fisica 7, 25 (1961).

Conforto, A. M. and N. Iucci. Cosmic Ray Events in November 1960, Nuovo Cimento, Series X, 21, 294 (1961).

Yasuhare, M. and H. Maeda. Geomagnetic Crochet of 15 November 1960, J. Atmos. Terres. Phys. 21, 278 (1961).

Campbell, W. S. and P. L. Hubert. The Ionospheric Disturbances of November 12 and 16, 1960, Can. J. Phys. 39, 614 (1961).

Lockwood, J. A. and M. A. Shea. Variations of the Cosmic Radiation in November 1960, J. Geophys. Res. 66, 3083 (1961).

Roederer, J. G. et al. Cosmic Ray Phenomena During the November 1960 Solar Disturbances, J. Geophys. Res. 66, 1603 (1961).

Steljes, J. F., H. Carmichael, and K. G. McCracken. Characteristics and Fine Structure of the Large Cosmic-Ray Fluctuations in November 1960, J. Geophys. Res. 66, 1363 (1961).

Wasson, J. T. Radioactivity Produced in Discoverer XVII by November 12, 1960, Solar Flare Protons, J. Geophys. Res. 66, 2659 (1961).

Carmichael, H. and J. F. Steljes. Review of Recent High Energy Solar Particle Events Including November 1960, Atomic Energy of Canada Ltd., AECL-1387 (CRGP--1056) October 30, 1961; also, Proceedings International Conference on Cosmic Rays and the Earth Storm, Kyoto, September 1961. II. Joint Sessions. Physical Society of Japan, Tokyo (1962), pp. 337-344.

Ney, E. P. and W. Stein. Solar Protons in November 1960, Proceedings of International Conference on Cosmic Rays and the Earth Storm, Kyoto, September 1961. II. Joint Sessions. Physical Society of Japan, Tokyo (1962), pp. 345-353.

Davis, L. R. and K. W. Ogilvie. Rocket Observations of Solar Protons during the November 1960 Events, 2, J. Geophys. Res. 67, 1711 (1962).



McCracken, K. G. The Cosmic Ray Flare Effect 2 - The Flare Effects of May 4, November 12, and November 15, 1960, J. Geophys. Res. 67, 435 (1962).

Ney, E. P. and W. A. Stein. Solar Protons, Alpha Particles, and Heavy Nuclei in November 1960, J. Geophys. Res. 67, 2087 (1962).

Ogilvie, K. W., D. A. Bryant, and L. R. Davis. Rocket Observations of Solar Protons during the November 1960 Events, 1, J. Geophys. Res. 67, 929 (1962).

Biswas, S., C. E. Fitchel, and D. E. Guss. Study of H, He, and Heavy Nuclei in November 12, 1960 Solar Cosmic Ray Event, Phys. Rev. 128, 2756 (1962).

Biswas, S. et al. Hydrogen, Helium, and Heavy Nuclei from the Solar Event on November 15, 1960, J. Geophys. Res. 68, 3109 (1963).

Ogilvie, K. W. and D. A. Bryant. Solar Proton Spectra in the Events of November 12 and 15, 1960, J. Geophys. Res. 69, 393 (1964).

Zirin, H. The Limb Flare of November 20, 1960. Astrophys. J. 140, 1216 (1964).

July 1961

Gaisauskas, V. and A. E. Covington. Radio and Corpuscular Radiation Associated with the Flare Surge on the Western Limb of the Sun on July 29, 1961, J. Geophys. Res. 67, 4119 (1962).

Machlum, B. and B. J. O'Brien. Solar Cosmic Rays of July 1961 and their Ionospheric Effects, J. Geophys. Res. 67, 3269 (1962).

Pieper, G. F. et al. Solar Protons and Magnetic Storms in July 1961, J. Geophys. Res. 67, 4959 (1962).

Ellison, M. A., S. M. P. McKenna, and J. H. Reid. Cosmic Ray Flares Associated with the 1961 July Event, Mon. Not. Roy. Astron. Soc. 124, 263 (1962).

Keppler, E., A. Ehmert, and G. Pfozter. Solar Proton Injections during the Period from July 12 to July 28, 1961 at Balloon Altitudes in the Auroral Zone, Space Res. 3, 676 (1963).



Hofmann, D. J. and J. R. Winckler. Simultaneous Balloon Observations at Ft. Churchill and Minneapolis during the Solar Cosmic Ray Events of July 1961, Space Res. 3, 662 (1963).

Kolomeets, E. V. Intensity Bursts of Cosmic Rays of July 18 and 20, 1961, in Section VII of Cosmic Rays (Results of Researches on the Program of the International Geophysical Year), S. N. Vernov and L. I. Dorman, eds., USSR Academy of Sciences (1963), p. 116.

Pieper, G. F., A. J. Zmuda, and C. O. Bostrom. Solar Protons and the Magnetic Storm of 13 July 1961, Space Res. 3, 649 (1963).

Krivsky, L. Cosmic Ray Flare of 1961 July 18 and Y-shaped Stage of Flares Development as Phase-Conditioning Ejection, Nuovo Cimento, Series X, 27, 1017 (1963).

Guss, D. E. and C. J. Waddington. Observations on the Solar Particle Events of July 1961, J. Geophys. Res. 68, 2619 (1963).

Hofman, D. J. and J. R. Winckler. Simultaneous Balloon Observations at Fort Churchill and Minneapolis during the Solar Cosmic Ray Events of July 1961, J. Geophys. Res. 68, 2067 (1963).

Bennett, S. M. Discussion of a Paper by D. J. Hofman and J. R. Winckler, "Simultaneous Balloon Observations at Fort Churchill and Minneapolis During the Solar Cosmic Ray Events of July 1961", J. Geophys. Res. 69, 2843 (1964).

Hofman, D. J. and J. R. Winckler. Reply, J. Geophys. Res. 69, 2845 (1964).

September 1961

Van Allen, J. A. et al. Coordinated Injun I--Explorer XII Observations of Solar Cosmic Rays, 28 September to 4 October 1961, presented at Explorer XII Symposium, Goddard Space Flight Center (1962).

Anderson, K. A. and J. R. Winckler. Solar X-Ray Burst on September 28, 1961, J. Geophys. Res. 67, 4108 (1962)

Bryant, D. A. et al. Explorer 12 Observations of Solar Cosmic Rays and Energetic Storm Particles after the Solar Flare of September 28, 1961, J. Geophys. Res. 67, 4963 (1962).



Hoffman, R. A., L. R. Davis, and J. M. Williamson. Protons of 0.1 to 5 Mev and Electrons of 20 kev at 12 Earth Radii during Sudden Commencement on September 30, 1961, J. Geophys. Res. 67, 5001 (1962).

Jean, A. G. and J. H. Crary. Very Low Frequency Phase Observations of the Ionospheric Effects of the Solar Flare of September 28, 1961, J. Geophys. Res. 67, 4903 (1962).

Kanellakos, D. P. and O. G. Villara. Ionospheric Disturbances Associated with the Solar Flare of September 28, 1961, J. Geophys. Res. 67, 2265 (1962).

Maxwell, A. Generation of Radio Waves and Acceleration of Particles by the Class 3 Solar Flare of 1961 September 28, Planet. Space Sci. 11, 897 (1963).

Dorman, L. I. and L. I. Miroshnichenko. The Solar Cosmic Ray Burst of 28 September 1961, Geomagnetism and Aeronomy 5, 377 (1965).

November 1961

Bates, H. F. Very Low Frequency Effects from the November 10, 1961 Polar Cap Absorption Event, J. Geophys. Res. 67, 2745 (1962).

February 1962

Zmuda, A. J., G. F. Pieper, and C. O. Bostrom. Solar Protons and Magnetic Storms in February 1962, J. Geophys. Res. 68, 1160 (1963).

September 1962

Rao, U. R. Study of Cosmic Ray Intensity Measured by Mariner 2 and by Ground Neutron Monitors During the Forbush Decrease of September 30, 1962, J. Geophys. Res. 70, 3765 (1965).

September 1963

Van Allen, J. A. Solar Cosmic-Ray Events January-September 1963, Trans. Am. Geophys. Union 45, 351 (1964).

Goedeke, A. D. and A. J. Masley. Observations in the Antarctic of Solar Cosmic Ray Events in 1962 and 1963, J. Geophys. Res. 69, 4166 (1964).

Moreton, G. E. H α Shock Wave and Winking Filaments with the Flare of 20 September 1963, Astron. J. 69, 145 (1964).

Lazuten, L. L. Solar Cosmic Rays Recorded in the Stratosphere in September 1963, Geomagnetism and Aeronomy 5, 563 (1965).

Cline, T. C. et al. Explorer 14 Observations of Solar Proton Events and Interplanetary Particle Events in 1962 and 1963, Trans. Am. Geophys. Union 46, 129 (1965).



PREDICTION OF SOLAR FLARES

Dodson, H. W., E. R. Hedeman, and J. Chamberlin. Ejection of Hydrogen and Ionized Calcium Atoms with High Velocity at the Time of Solar Flares, Astrophys. J. 117, 66 (1953).

Billings, D. E. Photospheric Magnetic Fields Preceding Solar Activity, Astrophys. J. 120, 184 (1954).

Warwick, C. S. Some Characteristics of Solar Flares, Astrophys. J. 120, 237 (1954).

Howard, R. An Analysis of the Spectra of Sunspots, Astron. J. 62, 143 (1957).

Warwick, J. W. Flare-Connected Prominences, Astron. J. 125, 811 (1957).

Granger, C. W. J. A Statistical Model for Sunspot Activity, Astrophys. J. 126, 152 (1957).

Billings, D. E. H α Profiles and Opacity of Small Flares, Astron. J. 63, 303 (1958).

Dodson, H. W. and E. R. Hedeman, Crochet-Associated Flares, Astron. J. 63, 304 (1958).

Dodson, H. W. and E. R. Hedeman. Crochet-Associated Flares, Astrophys. J. 128, 636 (1958).

Evans, J. W. Flare-Associated Magnetic Activity in the Sun, Astron. J. 64, 330 (1959).

Warwick, C. and M. Wood. A Study of Limb Flares and Associated Events, Astrophys. J. 129, 801 (1959).

Leighton, R. B. Observations of Solar Magnetic Fields in Plage Regions, Astrophys. J. 130, 366 (1959).

Denisse, J. F., A. Boischot, and M. Pick-Gutmann. Properties of Chromospheric Eruptions Associated with Production of Cosmic Rays by the Sun, Space Res. 1, 637 (1960).

Dodson, H. W. and E. R. Hedeman. Survey of Number of Solar Flares Observed During the International Geophysical Year, J. Geophys. Res. 65, 123 (1960).



Dodson, H. W. and E. R. Hedeman. Survey of Number of Flares Observed During the IGY, Astron. J. 65, 51 (1960).

Minnis, C. M. An Estimate of the Peak Sunspot Number in 1968, J. Atmos. Terres. Phys. 20, 94 (1961).

Biller, W. Solar Ray Storm Prediction Improved, Missiles and Rockets 9, 30 (August 7, 1961).

Sandstrom, A. E. Prediction of Cosmic Ray Storms from Geomagnetic Data, Astronautica Acta 7, 141 (1961).

Westfall, W. D. Prediction of VLF Diurnal Phase Changes and Solar Flare Effect, J. Geophys. Res. 66, 2733 (1961).

Westfall, W. D. Comments on Paper, Prediction of VLF Diurnal Phase Changes and Solar Flare Effect, J. Geophys. Res. 67, 916 (1962).

Westfall, W. D. Author's Reply to the Preceding Discussion, J. Geophys. Res. 67, 918 (1962).

Acton, L. Effect of Intermittent Operation of Flare Patrols on Observed Flare Frequency, Astron. J. 67, 109 (1962).

Haurwitz, M. W. Delay Time from Flare to SC Storm as a Function of Pre-Existing Geomagnetic Conditions, Astron. J. 67, 116 (1962).

Warwick, C., M. W. Haurwitz, and C. E. Hoff. Investigation and Elimination of Systematic Errors in Measures of Solar Flares, Astron. J. 67, 123 (1962).

Zirin, H. Physical Conditions in Limb Flares and Active Prominences. V - Excitation and Ionization of Helium and Metals, Astrophys. J. 135, 521 (1962).

Malitson, H. H. Predicting Large Solar Cosmic Ray Events, Astronautics and Aerospace Engineering 1, 70 (March 1963).

Hakura, Y. and J. V. Lincoln. A Suggestion for Improving Forecasts of Geomagnetic Storms, J. Geophys. Res. 68, 1163 (1963).

Kamphouse, J. L. Correlation of Neutron Monitor Pressure Coefficient and the Solar Cycle, J. Geophys. Res. 68, 5609 (1963).

Ramsey, H. E. and S. F. Smith. Pre-Flare Filament Characteristics, Astron. J. 68, 290 (1963).



Greatrix, G. R. On the Statistical Relations Between Flare Intensity and Sunspots, Mon. Not. Roy. Astron. Soc. 126, 123 (1963).

Byrne, F. N., M. A. Ellison, and J. H. Reid. A Survey of Solar Flare Phenomena, Space Science Review 11, 319 (1964).

Guss, D. E. Distribution in Heliographic Longitude of Flares Which Produce Energetic Solar Particles, Phys. Rev. Ltrs. 13, 363 (1964).

Jonah, F. D. Solar Activity Catalog for the Nineteenth Solar Cycle, Trans. Am. Geophys. Union 45, 626 (1964).

Masley, A. J., G. W. Adams, and A. D. Goedeke. Solar Cosmic Ray Event Characteristics Near Solar Minimum, Trans. Am. Geophys. Union 45, 626 (1964).

Gregory, J. B. and R. E. Newdick. Twenty-seven Day Recurrence of Solar Protons, J. Geophys. Res. 69, 2383 (1964).

Dodson, H. W. and E. R. Hedeman. An Unexpected Effect in Solar Cosmic Ray Data Related to 29.5 Days, J. Geophys. Res. 69, 3965 (1964).

Dodson, H. W. and E. R. Hedeman. Solar Minimum and the International Years of the Quiet Sun, Science 43, 237 (1964).

Gopasiuk, S. I. et al. On the Forecasting of Solar Flares, International Astronautical Congress, 13th, Varna, Bulgaria, September 1962, Proceedings, Vol. 1 (1964), pp. 63-69.

Morney, A. E. Study of the Application of Perceptrons for the Prediction of Large Solar Flares, Prep. II Monthly Letter Report No. 2, June 1964, Lockheed Aircraft Corp. (July 15, 1964).

Grebinskii, A. S. and A. P. Molchanov. Some Characteristics of Radiation Sources on the Sun Associated with Groups of Spots, Geomagnetizm i Aeronomiia, 4, 635 (1964).

Weddell, J. B. Prediction of Probability of Occurrence of Solar Flares, Astronautica Acta 10, 339 (1964).

Devaney, R. A., R. C. Blanchard, and J. Andary. Time Varying Solar Cycle Protons Program Manual, NASA-TM-X-55134; X-642-64-278, #N 65-18257 (Oct. 1964).



- Firor, J. and C. Lilliequist. Solar Flares and Prediction, Southwest Research Institute and USAF, International Symposium on Bioastronautics and the Exploration of Space, 3rd, San Antonio, Texas, (Nov. 16-18, 1964).
- Blizard, J. B. Prediction of Solar Flares Months in Advance, Astron. J. 70, 667 (1965).
- Blizard, J. B. and B. C. Epstein. The Persistence and Rotation Rate of Flare Centers, Denver Research Institute, (29 Dec. 1965).
- Blizard, J. B. A Long Range Solar Flare Prediction Technique, Denver Research Institute, (1965).
- McIntosh, P. S. On the Association of Solar Flares with Sunspots, Astron. J. 70, 673 (1965).
- McKenna, S. M. P. Positions of Certain Recurrent Flares Relative to Sunspots and Features of the Calcium Network, Astron. J. 70, 684 (1965).
- Shipiro, R. Comparisons of Power Spectrums of Artificial Time Series with Spectrum of a Solar Plage Index, J. Geophys. Res. 70, 3581 (1965).
- Callendar, R. H., J. R. Manzano, and J. R. Winckler. The Response of High Altitude Ionization Chambers During the 1954-1965 Solar Cycle, J. Geophys. Res. 70, 3189 (1965).
- Warwick, C. S. Solar Flare Frequency and Observing Time Patterns, Astrophys. J. 142, 767 (1965).
- Dodson, H. W., E. R. Hedeman, and F. L. Stewart. Solar Activity During the First 14 Months of the International Years of the Quiet Sun, Science 148, 1328 (1965).
- Theiss, C. M. and A. E. Murray. Study of the Application of Perceptrons for Prediction of Solar Flares, Solar Flare Forecasting with Recognizing Automation, Final Report, Phase II, Lockheed Aircraft Corp. (February 1965).
- Sand, R. C., Jr. Solar Flare Hazards: Predictability, General Electric Co., Philadelphia, Missile and Space Division (July 1965).
- Michaels, D. Forecasting Solar Flares, Foreign Sci. Bull. 1, No. 8, 18 (August 1965).



Vitinskii, Y. I. Solar Activity Forecasting, USSR Academy of Sciences, Leningrad (1962) - Translated by Israel Program for Scientific Translations, Jerusalem, (1965).

ORIGIN OF SOLAR FLARES

Richardson, R. S. Characteristics of Solar Flares, Astrophys. J. 114, 356 (1951).

Billings, D. E. and W. O. Roberts. Isophotal Photometry of a Solar Flare, Astrophys. J. 118, 429 (1953).

Larmore, L. A Study of the Motions of Solar Prominences, Astrophys. J. 118, 436 (1953).

Dolder, F. P., W. O. Roberts, and D. E. Billings. Solar Flares and the Yellow Coronal Line, Astrophys. J. 119, 120 (1954).

Goldberg, L., H. W. Dodson, and E. A. Muller. The Width of H α in Solar Flares, Astrophys. J. 120, 83 (1954).

Menzel, D. H., D. Layzern, and M. Krook. The Origin and Evolution of Sunspots, Astron. J. 60, 171 (1955).

Roberts, W. O. and D. E. Billings. A Hypothesis for the Origin of Solar Flares, Astron. J. 60, 176 (1955).

Warwick, J. J. Heights of Solar Flares, Astrophys. J. 121, 376 (1955).

Dodson, H. W., E. R. Hedeman, and R. R. McMath. Photometry of Solar Flares, Astrophys. J. 123, 186 (1956).

Warwick, J. W. Flare-Connected Prominences, Astrophys. J. 125, 811 (1957).

Gold, T. and F. Hoyle. On the Origin of Solar Flares, Mon. Not. Roy. Astron. Soc. 128, 89 (1960).



Dodson, H. W. Observation of Loop-Type Prominences in Projection Against the Disk at the Time of Certain Solar Flares, Proc. Nat. Acad. Science 47, (7), 901 (1961).

Parker, E. N. The Solar Flare Phenomenon and the Theory of Reconnection and Annihilation of Magnetic Fields, Astrophys. J. 138, 310 (1963).

Wentzel, D. G. Solar Flares Caused by the Skin Effect in Twisted Magnetic Fields, Astrophys. J. 140, 1563 (1964).

Severny, A. B. Solar Flares, Ann. Rev. Astron. Astrophys. 2, (1964).

Ellison, M. A. and J. H. Reid. Solar Flares, Research in Geophysics, H. Odishaw, ed., Vol. 1, Sun, Atmosphere, and Space. MIT Press, Cambridge, Mass. (1964), pp. 43-63.

Glaser, H. Solar Flares, Naval Research Reviews 18, 1 (March 1965).

Sturrock, P. A. and B. Coppi. A New Model of Solar Flares, Astrophys. J. 143, 3 (1966).

SOLAR RADIO BURSTS

Broxon, J. W. Further Investigation of Frequency of Occurrence of Small Cosmic Ray Bursts, Phys. Rev. 72, 525 (1947).

Dodson, H. W., E. R. Hedeman, and L. Owren. Solar Flares and Associated 200 Mc/Sec Radiation, Astrophys. J. 118, 169 (1953).

Dodson, H. W., E. R. Hedeman, and A. E. Covington. Solar Flares and Associated 2800 Mc/Sec (10.7 cm) Radiation, Astrophys. J. 119, 541 (1954).

Hanssen, E. T. On the Correlation Between Radio-Frequency Radiation from the Sun and Solar Activity, Astrophys. J. 121, 367 (1955).

De Jager, C. Optical and Radiophenomena Associated with a Solar Activity Center, Nuovo Cimento, Series X, 13, 284 (1959).

Edelson, S. et al. Time Relations Between Centimeter Wavelength Bursts and Solar H α Flares, Astron. J. 64, 330 (1959).

Hachenberg, O. and A. Kruger. The Correlation of Bursts of Solar Radio Emission in the Centimetre Range with Flares and Sudden Ionospheric Disturbances, J. Atmos. Terres. Phys. 17, 20 (1959).



Bachelet, F., A. M. Conforto, and N. Iucci. Solar Flares with Type IV Radioburst and Transient Phenomena of Cosmic Rays, Space Res. 1, 662 (1960).

Thompson, A. R. and A. Maxwell. Solar Radio Bursts and Low-Energy Cosmic-Rays, Nature 185, 89 (1960).

Bachelet, F. et al. Cosmic Ray and Geomagnetic Disturbances from July 1957 to July 1958 III - The Correlation with Solar Radio Bursts, Nuovo Cimento, Series X, 21, 648 (1961).

Suburai, K. and H. Maeda. A Relation between Solar Radio Emission and Low-Energy Solar Cosmic Rays, J. Geophys. Res. 66, 1966 (1961).

Smith, E. P. Type V Bursts and Their Association with Flares, and Addenda to the Flare Association of Type I Bursts, J. Geophys. Res. 67, 3797 (1962).

Malville, J. M. The Association of Type III Bursts and Solar Flares, Astrophys. J. 135, 834 (1962).

Stein, W. A. and E. P. Ney. Continuum Electromagnetic Radiation from Solar Flares, J. Geophys. Res. 68, 65 (1963).

Chan, K. L. and O. G. Villard. Sudden Frequency Deviations Induced by Solar Flares, J. Geophys. Res. 68, 3197 (1963).

Das Gupta, M. K. and D. Basu. Solar Radio Bursts in Relation to Flare Importances, Astrophys. J. 137, 997 (1963).

Haurwitz, M. W. Solar Flares with Associated Active Dark Filaments and Their Relation to 2800 Mc/s Radio Bursts, Astrophys. J. 140, 1236 (1964).

Mitra, S. N. A Radio Method of Detecting Solar Flares, J. Atmos. Terres. Phys. 26, 375 (1964).

Zancha, H. and Z. Kakinuma. The Relation Between the Spectrum of Slowly Varying Component of Solar Radio Emission and Solar Proton Event, Report on Ionosphere and Space Research in Japan, 18, No. 1, 32 (1964).

Smerd, S. F. Solar Radio Emissions, Research in Geophysics, H. Odishaw, ed., Vol. 1, Sun, Upper Atmosphere, and Space. MIT Press, Cambridge, Mass. (1964), pp. 65-97.



Acton, L. W. Some Relationships Between Short-Wave Fadeouts, Magnetic Crochets, and Solar Flares, J. Geophys. Res. 66, 3060 (1961).

Corkhill, P. J., R. A. Hoffman, and R. J. Lockhart. Measurement of Solar Flare-Produced Cosmic Radiation, Aerospace Medical Division, Aeromedical Research Lab., Holloman AFB, N. Mex. (October 1962).

McCracken, K. G. The Cosmic-Ray Flare Effect. 1, Some New Methods of Analysis, J. Geophys. Res. 67, 423 (1962).

Vogt, R. Primary Cosmic-Ray and Solar Protons, Phys. Rev. 125, 366 (1962).

Dubov, E. E. A Possible Mechanism of Radiation Emission in Chromospheric Flares, Soviet Physics - Doklady 8, 543 (1963).

Jaggi, R. K. The Role of Plasma Instabilities in the Origin of Solar Flares, J. Geophys. Res. 68, 4429 (1963).

Zirin, H. Coronal Emission from Solar Flares, Astron. J. 68, 82 (1963).

Meyer, P. and R. Vogt. Primary Cosmic Ray and Solar Protons II, Phys. Rev. 129, 2275 (1963).

Malville, J. M. The Association of Energetic Particles with Solar Flares, AAS-NASA Symposium on the Physics of Solar Flares, NASA-SP-50, pp. 257 (1963).

McDonald, F. B. Solar Cosmic Rays, presented at 31st Annual Meeting, Institute of Aerospace Sciences, New York (1963).

Reitz, D. Solar Cosmic Ray Events and Their Interpretation, Part I Introduction, Martin Marietta Corp. Report (May 1963).

Reitz, D. Solar Cosmic Ray Events and Their Interpretation, Part II Measurements, Martin Marietta Corp. Report (July 1963).

Parker, E. N. Coronal Expansion and Solar Corpuscular Radiation, Research in Geophysics, H. Odishaw, ed., Vol. 1, Sun, Upper Atmosphere, and Space. MIT Press, Cambridge, Mass. (1964), pp. 99-114.

Axford, W. I. and E. C. Ray. A Model for Solar Proton Events, Trans. Am. Geophys. Union 45, 79 (1964).

Wentzel, D. G. Formation of Solar Flares Without Magnetic Neutral Points, Astron. J. 69, 145 (1964).



Fletcher, J. D. Solar Radio Emission as a Criterion for Solar Proton Event Warning, AIAA Journal 2, 2193 (1964).

Yoshida, S. Solar Cosmic Rays Associated with Meter Type IV Radio Bursts, Geophys. Res. 70, 2065 (1965).

Radicella, S. M. and M. A. Sette. Effects of Solar Radio Bursts on Riometer Records at 30 Mc/s, J. Geophys. Res. 70, 5970 (1965).

McKenna, S. M. P. A Correlation Between Optical and Radio Aspects of a Class 3 Flare, Mon. Not. Roy. Astron. Soc. 129, 437 (1965).

SOURCE OF SOLAR FLARE PARTICLES

Forbush, S. E. Three Unusual Cosmic Ray Increases Possibly Due to Charged Particles from the Sun, Phys. Rev. 70, 771 (1946).

Kiepenheuer, K. O. On the Origin of the Cosmic Radiation from the Sun, Phys. Rev. 78, 809 (1950).

Lord, J. J., A. W. Elston, and M. Schein. A Large Fluctuation in the Rate of Production of Nuclear Disintegrations Following a Solar Flare, Phys. Rev. 79, 540 (1950).

Pomerantz, M. A. An Increase of the Primary Cosmic-Ray Intensity Following a Solar Flare, Phys. Rev. 81, 731 (1951).

Simpson, J. A., W. Fonger, and L. Wilcox. Solar Component of Primary Cosmic Radiation, Phys. Rev. 85, 366 and 720 (1952).

Pomerantz, M. A. and G. W. McClure, On the Enhanced Primary Cosmic Ray Intensity Associated with Solar Disturbances, Phys. Rev. 87, 240 (1952).

Dodson, H. W., E. R. Hedeman, and J. Chamberlin. Ejection of Hydrogen and Ionized Calcium Atoms with High Velocity at the Time of Solar Flares, Astrophys. J. 117, 66, (1953).

Firor, J. Cosmic Radiation Intensity-Time Variations and Their Origin IV - Increases Associated with Solar Flares, Phys. Rev. 94, 1017 (1954).

Sekido, J. Cosmic Rays and Corpuscles from the Sun, Reports on Ionosphere Research in Japan 9, 181 (1955).

Ehmert, A. and G. Pfozter. A New Outburst of Solar Cosmic Rays, Z. Naturforsch 11a, 322 (1956).



Parker, E. N. Acceleration of Cosmic Rays in Solar Flares, Phys. Rev. 107, 830 (1957).

Boischot, A. and J. F. Denisse. Les Émissions de Type IV et l'Origine des Rayons Cosmiques Associés aux Éruptions Chromosphériques, Comptes Rend. 245, 2194 (1957).

Corrigan, J. J., S. F. Singer, and M. J. Swetnik. Cosmic-Ray Increases Produced by Small Solar Flares, Phys. Rev. Ltrs. 1, 104 (1958).

Sweet, P. A. The Production of High Energy Particles in Solar Flares, Nuovo Cimento, Series X, Supplement, 8, 188 (1958).

Cocconi, G. et al. The Cosmic Ray Flare Effect, Nuovo Cimento, Series X, Supplement 8, 161 (1958).

Katzman, J. A Cosmic Ray Increase Related to Solar Activity, Can. J. Phys. 36, 807 (1958).

McCracken, K. G. Emission of White Light and Cosmic Radiation by a Solar Flare, Nuovo Cimento, Series X, 13, 1081 (1959).

Ward, W. H. On the Origin of Terrestrial Particles from Solar Flares, J. Atmos. Terres. Phys. 14, 296 (1959).

Katzman, J. The Sun as a Source of Cosmic Rays of Intermediate Energies, Can. J. Phys. 37, 1207 (1959).

Towle, L. C. and J. A. Lockwood. Cosmic Ray Increases Associated with Solar Flares, Phys. Rev. 113, 641 (1959).

Denisse, J. F., A. Boischot, and M. Pick-Gutmann. Properties of Chromospheric Emissions Associated with Production of Cosmic Rays by the Sun, Space Res. 1, 637 (1960).

Charakhchian, A. N., V. F. Tulinov, and T. N. Charakhchian. Cosmic Rays Emitted by the Sun, Space Res. 1, 649 (1960).

Fichtel, C. E. and D. E. Guss. Heavy Nuclei in Solar Cosmic Rays, Phys. Rev. Ltrs. 6, 495 (1961).

Carmichael, H. et al. Cosmic Ray Neutron Increase from a Flare on the Far Side of the Sun, Phys. Rev. Ltrs. 6, 49 (1961).

Tulinov, V. F. and T. N. Charakhchian. Cosmic Rays from the Sun, Soviet Phys. - JETP 12, 179 (1961).



Bruzek, A. Optical Characteristics of Cosmic-Ray and Proton Flares, J. Geophys. Res. 69, 2386 (1964).

Reid, G. C. A Diffusive Model for the Initial Phase of a Solar Proton Event, J. Geophys. Res. 69, 2659 (1964).

Waddington, C. J. and P. S. Freier, Relative Abundances of Energetic Hydrogen Isotopes Produced in Solar Flares, Phys. Rev. 136, 1535 (1964).

Tandburg-Hanssen, E. Solar Flares and the Associated Ejection of Particles, in Virginia Polytechnic Inst. Conference on Artificial Satellites, Part B, (August 1964).

Avignon, Y., M. J. Martines, and M. Pide. The Identification of Chromospheric Eruption Classes Associated with Ray Emissions and Radioelectric Activity, Ann. d' Astrophys. 27, 23 (1964).

Biswas, S. and C. E. Fichtel. Composition of Solar Cosmic Rays, Space Science Revs. 4, 709 (1965).

Bryant, D. A. et al. Continual Acceleration of Solar Protons in the Mev Range, Phys. Rev. Ltrs. 14, 481 (1965).

Lingenfelter, R. E. et al. High Energy Solar Neutrons 2 - Flux at the Earth, J. Geophys. Res. 70, 4087 (1965).

Sturrock, P. A. and B. Coppi. A New Model of Solar Flares, Astron. J. 70, 331 (1965).

Weddell, J. B. Production of Energetic Solar Proton Events by Hydro-magnetic Shocks, Planet. Space Sci. 13, 785 (1965).

Wentzel, D. G. Fermi Acceleration of Solar Cosmic Rays, J. Geophys. Res. 70, 2716 (1965).

Lingenfelter, R. E. et al. High Energy Solar Neutrons 1 - Production in Flares, J. Geophys. Res. 70, 4077 (1965).

Miroshnichenko, L. I. Cosmic Rays from the Sun, NASA TT-F-9648; ST-CR-SP-10299, #N 65-18183, (4 March 1965).

Head, R. M. Solar Proton Event Centers of the Chromosphere, COSPAR, International Space Science Symposium, 6th, Buenos Aires, Argentina, 13-19 May 1965.



PROPAGATION OF SOLAR FLARE PARTICLES

Lüst, R. and J. A. Simpson. Initial Stages in the Propagation of Cosmic Rays Produced by Solar Flares, Phys. Rev. 108, 1563 (1957).

Simpson, J. A. Solar Flare Cosmic Rays and Their Propagation, Nuovo Cimento, Series X, Supplement 8, 133 (1958).

Singer, S. F. A Model for Solar Flare Increases of Cosmic Rays Based on Hydromagnetic Turbulence, Nuovo Cimento, Series X, Supplement 8, 197 (1958).

Ogayashi, T. and Y. Kakura. Propagation of Solar Cosmic Rays Through Interplanetary Magnetic Field, J. Geophys. Res. 65, 3143 (1960).

McCracken, K. G. The Cosmic-Ray Flare Effect. 3, Deductions Regarding the Interplanetary Magnetic Field, J. Geophys. Res. 67, 447 (1962).

Beard, D. B. The Interaction of the Terrestrial Magnetic Field with the Solar Corpuscular Radiation 2, Second Order Approximation, J. Geophys. Res. 67, 477 (1962).

Warwick, C. S. Propagation of Solar Particles and the Interplanetary Magnetic Field, J. Geophys. Res. 67, 477 (1962).

Greenstadt, E. W. A Comparison of Solar Flare Incidence with Magnetic Transients Observed in the Near-by Interplanetary Region by Pioneer 5, J. Geophys. Res. 67, 3299 (1962).

Fireman, E. L. Density of the Solar Flare Plasma, J. Geophys. Res. 67, 4890 (1962).

Warwick, C. S. Solar Particles and Interplanetary Fields, presented at American Astronautical Society, 9th Annual Meeting, Los Angeles (15-17 January 1963).

Bryant, D. A. et al. New Evidence for Long Lived Solar Streams in Interplanetary Space, Phys. Rev. Ltrs. 11, 144 (1963).

Yoshida, S. et al. Propagation of Solar Particles Deduced from Measurements with Explorer VII Space Res. 3, 608 (1963).

McCracken, K. G. Solar Cosmic Ray Propagation Characteristics, International Conference on Cosmic Rays, 8th, Jaipur, India, (1963 December 2-14), Proceedings. Vol. 1, Solar Particles and Sun-Earth Relations, pp 83-94.



Krymskii, G. F. and A. I. Kuzmin. Solar Cosmic Rays and Interplanetary Magnetic Field, International Conference on Cosmic Rays, 8th, Jaipur, India, (2-14 December 1963), Proceedings. Vol. 1, Solar Particles and Sun-Earth Relations, pp 226-229.

Obayashi, S. Some Notes on Low Energy Solar Cosmic Rays and Interplanetary Magnetic Fields, Space Science 3, 624 (1963).

Obayashi, T. The Streaming of Solar Flare Particles and Plasma in Interplanetary Space, Space Science Rev. 3, 79 (1964).

Roederer, J. G. Generation, Propagation and Detection of Relativistic Solar Particles, Space Science Rev. 3, 487 (1964).

Obayashi, T. Streaming of Solar Particles between Sun and Earth, Planet. Space Sci. 12, 463 (1964).

Dorman, L. I. and L. I. Miroshnichenko. Energy Dependence of the Coefficient of Diffusion of Solar Cosmic Rays, Geomagnetizm i Aeronomiia 4, 127 (1964).

Krymskii, G. F. and A. I. Kuzmin. Solar Cosmic Rays and the Interplanetary Magnetic Field, Bull. Acad. Sci. USSR, Physical Series, 28, 1896 (1964).

Dorman, L. I. and Miroshnichenko. Concerning the Character and Propagation of Solar Cosmic Rays in Interplanetary Space, Bull. Acad. Sci. USSR, Physical Series 28, 1870 (1964).

Boldt, Elihu. Do Solar Flare Proton Events Measure Geometry of a Diffusion Process, NASA-TM-X-55161, #N 65-18275, (Dec. 1964).

Yates, G. K. Solar Flare High Energy Alpha Particles and Their Storage in Interplanetary Space, J. Geophys. Res. 69, 3077 (1964).

Waddington, C. J. and P. S. Freier. Comment on a Paper by G. K. Yates, Solar Flare High Energy Alpha Particles and Their Storage in Interplanetary Space, J. Geophys. Res. 70, 230 (1965).

Yates, G. K. Reply, J. Geophys. Res. 70, 232 (1965).

Fibich, M. and P. B. Abraham. On the Propagation and Diffusion of Solar Protons, J. Geophys. Res. 70, 2475 (1965).

Krimigis, S. M. Interplanetary Diffusion Model for the Time Behavior of Intensity in a Solar Cosmic Ray Event, J. Geophys. Res. 70, 2943 (1965).



Bennett, S. M. Solar Cosmic Rays in a Regular Interplanetary Magnetic Field, Scientific Report No. 19, Avco Corporation, (27 Jan 1965).

Yashida, S. and S. I. Akasofu. A Study of the Propagation of Solar Particles in Interplanetary Space - The Center Limb Effect of Magnitude of Cosmic-Ray Storms and of Geomagnetic Storms, Planet. Space Sci. 13, 435 (1965).

Kunitamo, S. Propagation of Solar Protons and the Interplanetary Magnetic Field, Planet. Space Sci. 13, 745 (1965).

Kriusky, L. Flight Time of Solar Cosmic and Proton Radiation to the Earth, Astron. Inst. Czech. Bulletin 16, No. 1, 27 (1965).

SPATIAL DISTRIBUTIONS OF SOLAR FLARE PARTICLES

Arnoldy, R. L., R. A. Hoffman, and J. R. Winckler. Solar Cosmic Rays and Soft Radiation Observed at 5,000,000 Kilometers from Earth, J. Geophys. Res. 65, 3004 (1960).

Bryant, D. A., T. L. Cline, et al. Results From the Goddard Cosmic Ray Experiments on Explorer XII, Presented at Explorer XII Symposium, Goddard Space Flight Center, Greenbelt, Md. (1962).

Dorman, L. I. and L. I. Miroshnichenko. Character of the Spatial - Temporal Distribution of Solar Cosmic Rays in Interplanetary Space, Geomagnetizm i Aeronomiia, 4, 730 (1964).

Dorman, L. I. and L. I. Miroshnichenko. The Nature of the Space-Time Distribution of Solar Cosmic Rays in Interplanetary Space, Geomagnetizm i Aeronomiia 4, 940 (1964).

Van Allen, J. A., L. A. Frank, and D. Venkatesan. Small Solar Cosmic Ray Events Observed with Mariner 2, Trans. Am. Geophys. Union 45, 80 (1964).

Anderson, H. R. Energetic Particles Measured Near Venus by Mariner 2, J. Geophys. Res. 69, 2651 (1964).

Stazer, A. K. A Model Anisotropic Solar Proton Event, American Inst. of Aeronautics and Astronautics, Aerospace Sciences Meeting, New York (20-22 January 1964).

Greenstadt, E. W. Interplanetary Magnetic Effects of Solar Flares - Explorer 18 and Pioneer 5, J. Geophys. Res. 70, 5451 (1965).



COSMIC RAY EFFECTS

Adams, N. A Temporary Increase in the Neutron Component of Cosmic Rays, Phil. Mag. 41, 503 (1950).

Broxon, J. W. and H. W. Boehmer. Cosmic Ray Intensity Following a Solar Flare, Phys. Rev. 78, 411 (1950).

Treiman, S. B. Solar Magnetic Moment and Cosmic Ray Effects Associated with Solar Flares, Phys. Rev. 94, 1029 (1954).

Morrison, P. Solar Connected Variations of Cosmic Rays, Phys. Rev. 95, 646 (1954).

Trumpy, B. On the Correlation Between the Intensity of Different Components of Cosmic Rays, The Meteorological Conditions and Magnetic Storms, Univ. Bergen Arbok, Paper 2, 19 pp. (1954).

Morrison, P. Solar Origin of Cosmic Ray Time Variations, Phys. Rev. 101, 1397 (1956).

Pomerantz, M. A. Cosmic Ray Observations at Very High Altitudes During Periods of Intense Solar Activity, Phys. Rev. 102, 870 (1956).

Parker, E. N. Modulation of Primary Cosmic-Ray Intensity, Phys. Rev. 103, 1518 (1956).

Barton, J. C. and Stockhausen, J. H. Increase of Cosmic Ray Intensity in Jamaica Due to a Solar Flare, J. Atmos. Terres. Phys. 7, 161 (1956).

Dorman, L. I. Cosmic Ray Variations, Moscow State Publishing House for Technical and Theoretical Literature, 1957: translation prepared by Technical Documents Liaison Office, Wright-Patterson Air Force Base (1958).

Neher, N. V. and S. E. Forbush. Correlation of Cosmic Ray Intensity and Solar Activity, Phys. Rev. Ltrs. 1, 173 (1958).

Ramaswamy, G. and S. D. Chattejee. Cosmic Ray Intensity Variations at Sea Level During Magnetic Storm Periods, Can. J. Phys. 36, 635 (1958).

Fenton, A. G., K. B. Fenton, and D. C. Rose. The Variations of Sea Level Cosmic Ray Intensity between 1954 and 1957, Can. J. Phys. 36, 824 (1958).

Webber, W. R. and J. J. Quenby. On the Derivation of Cosmic Ray Specific Yield Functions, Phil. Mag. 4, 654 (1959).

Kolomeets, E. V. Small Effects of Solar Flares and the Energy Spectrum of Primary Variation of Cosmic Rays, Soviet Physics-JETP 9, 960 (1959).

Fenton, A. G. et al. The Onset Times of Forbush-type Cosmic Ray Intensity Decreases, Can. J. Phys. 37, 970 (1959).



McDonald, F. B. and W. R. Webber. The Variation of the Cosmic Ray Intensity During a Solar Cycle, Space Res. 1, 968 (1960).

Charakhch'yan, A. N., V. F. Tulinov, and T. N. Charakhch'yan. Large Cosmic-Ray Intensity Fluctuations in the Stratosphere, Soviet Physics-JETP 11, 742 (1960).

Tandberg-Hanssen, E. Excitation of Metal Lines in Flares and Active Prominences, Astron. J. 67, 588 (1962).

Wilson, G. B. and C. P. Hehra. A Statistical Analysis of Cosmic Ray Intensity Changes Associated with Solar Flares, J. Geophys. Res. 67, 3707 (1962).

Webber, W. R. Time Variations of Low Energy Cosmic Rays During the Recent Solar Cycle, in Progress in Elementary Particle and Cosmic Ray Physics, ed. by J. G. Wilson and S. A. Wouthuysen, North Holland Publ. Co., Amsterdam, Vol. 6 (1962).

Simpson, J. A. Recent Investigations of the Low Energy Cosmic and Solar Particle Radiations, presented at the Conference on Cosmic Rays in Interplanetary Space, Pontifical Academy of Science, Vatican City, (October 1962).

Wilson, B. G. and C. P. Nehra. Changes in Cosmic Radiation Intensity Associated with Solar Flares, J. Geophys. Res. 68, 3305 (1963).

Dorman, L. I. and E. V. Kolomeets. The Effect of Small Solar Flares in the Hard Component of Cosmic Rays, (Results of Researches on the Programme of the International Geophysical Year. Section VII), Trans. Acad. Sci. USSR, p. 139 (1963).

Kolomeets, E. V. The Nature of the Solar Flare Effect in the Ionizing Component of Cosmic Rays, Moscow, Izdatel'stvo Akademii Nauk SSSR (1963), pp. 122-125. (in Russian) (1963).

Meyer, P. Cosmic Radiation and Solar Modulation, Research in Geophysics, H. Odishaw, ed., Vol. 1, Sun, Upper Atmosphere, and Space. MIT Press, Cambridge, Mass. (1964).

Fretter, W. B., R. R. Brown, and K. A. Anderson. Research in High Energy Particle Physics and Research on Cosmic Rays and Solar Particle Emissions, University of California, Berkeley, Final Report (15 December 1964).

Forman, M. A. Neutron Monitor Mass Absorption Coefficients at Chicago and Climax During Solar Cycle 19 (1954-1963), J. Geophys. Res. 70, 2469 (1965).

McDonald, F. B. Review of Galactic and Solar Cosmic Rays, NASA Goddard Space Flight Center, Greenbelt, Md. (March 1965).



Morozov, V. S. et al. Biological Effect of Cosmic Radiation in the Event that Solar Flares Occur on the Earth to Moon Route in Model Experiment, Intern. Cosmic Res. 3, 241 (May 1965).

ENERGY SPECTRA OF SOLAR FLARE PARTICLES

Yataro, S. and Y. Teitiro. Absorption of Flare-Type Cosmic Ray, J. Phys. Soc. Japan 4, 353 (1949).

Robey, D. H. Radiation Shield Requirement for Two Large Solar Flares, Astronautica Acta 6, 206 (1960).

Syrovatskii, S. I. Spectrum of Galactic and Solar Cosmic Rays, Soviet Physics-JETP 13, 1257 (1961).

Charakhchyan, A. A., V. F. Tulinov, and T. N. Charakhchyan. Energy Spectrum and Time Dependence of the Intensity of Solar Cosmic Ray Photons, Soviet Physics-JETP 14, 530 (1962).

Bailey, D. K. Time Variations of the Energy Spectrum of Solar Cosmic Rays in Relation to the Radiation Hazard in Space, J. Geophys. Res. 67, 391 (1962).

Schatzman, E. Energy and Mass Spectra of Solar Cosmic Rays, Space Res. 3, 709 (1963).

Freier, P. S. and W. R. Webber. Exponential Rigidity Spectra for Solar Flare Cosmic Rays, Trans. Am. Geophys. Union 44, 72 (1963).

Freier, P. S. and W. R. Webber. Exponential Rigidity Spectra for Solar Flare Cosmic Rays, J. Geophys. Res. 68, 1605 (1963).

Freier, P. S. Emulsion Measurements of Solar Alpha Particles and Protons, J. Geophys. Res. 68, 1805 (1963).

Bryant, D. A. et al. Velocity Dependence and Source Spectra of Solar Proton Events, International Conference on Cosmic Rays, 8th, Jaipur, India, 2-14 December 1963, Proceedings. Vol 1, Solar Particles and Sun-Earth Relations, pp. 70-79.

Lai, D., G. Rajagopalan, and V. S. Venkatavaradan. Radioactivity Induced by Solar Particles, International Conference on Cosmic Rays, 8th, Jaipur, India, 2-14 December 1963, Proceedings. Vol. 1, Solar Particles and Sun-Earth Relations, pp. 99-134.

Schatzman, E. Energy and Mass Spectra of Solar Cosmic Rays, Institut d'Astrophysique, Paris, pp. 709-715 (1963).



Freier, P.S. and W.R. Webber. Radiation Hazard in Space from Solar Particles, Science 142, 1598 (1963).

Lingenfelter, R.E. and E.J. Flamm. Production of Carbon 14 by Solar Protons, J. Atmos. Sciences 21, 134 (1964).

Biswas, S. and C.E. Fichtel. Nuclear Composition and Rigidity Spectra of Solar Cosmic Rays, Astrophys. J. 139, 941 (1964).

Dorman, L.I. and L.I. Miroshnichenko. Energy Dependence of the Coefficient of Diffusion of Solar Cosmic Rays, Geomagnetism and Aeronomy 4, 127 (1964).

Tonaka, H. and Z. Kakinuma. The Relation Between the Spectrum of Slowly Varying Component of Solar Radio Emission and Solar Proton Event, Report of Ionosphere and Space Research in Japan, 18, No. 1, 32 (1964).

McDonald, F.B. Velocity Dependence and Source Spectra of Solar Proton Events, in Virginia Polytechnic Inst. Conference on Artificial Satellites, Part A, (August 1964).

Roederer, J.G. High Energy Solar Particle Events, in Research in Geophysics, Vol. I, Sun, Upper Atmosphere, and Space, Hugh Odishaw, ed., MIT Press, Cambridge, Mass. (1964), pp. 115-145.

Charackhch'ian, A.A. and T.N. Charackhch'ian. The Energy Spectrum of Protons and Alpha Particles of Cosmic-Ray Flares on the Sun, Geomagnetizm i Aeronomiia 5, 220 (1965).

Sakurai, K. On the Nuclear Abundance and Rigidity Spectra of Solar Cosmic Rays, Planet. Space Sci. 13, 867 (1965).

POLAR CAP ABSORPTIONS

Skain, C.A. and A.P. Mitra. Effects of Solar Flares on the Absorption of 18.3 Mc/sec Cosmic Noise, J. Atmos. Terres. Phys. 5, 316 (1954).

Kundu, M.R. and F.T. Haddock. A Relation Between Solar Radio Emission and Polar Cap Absorption of Cosmic Noise, Nature 186, 610 (1960).

Obayashi, T. and Y. Hakura. Solar Corpuscular Radiation and Polar Ionospheric Disturbances, J. Geophys. Res. 65, 3131 (1960).

Herzberg, L. The Possible Importance of Nitric Oxide Formation During Polar-Cap Ionospheric Absorption Events, J. Geophys. Res. 65, 3505 (1960).



Jelly, D.H. Compiled Data for Polar Cap Absorption Events Observed at Churchill, June 1957 to June 1960, Canadian Defence Research Telecommunications Establishment Report No. 1062 (1961).

Reid, G.C. and H.O. Leinbach. Morphology and Interpretation of the Great Polar Cap Absorption Events of May and July 1959, J. Atmos. Terres. Phys. 23, 216 (1961).

Leinbach, H. Some Observations of Daytime Recoveries During Polar Cap Absorption Events, Arkiv for Geofysik 3, 427 (1961).

Ortner, J., H. Leinbach, and M. Sugiura. The Geomagnetic Storm Effect on Polar Cap Absorption, Arkiv for Geofysik 3, 429 (1961).

Reid, G.C. A Study of the Enhanced Ionization Produced by Solar Protons During a Polar Cap Absorption Event, J. Geophys. Res. 66, 4071 (1961).

Gregory, J.B. Ionospheric Reflections Below 50 Kilometers During Polar Cap Absorption Events, J. Geophys. Res. 66, 2575 (1961).

Rourke, G.F. Small-Scale Polar-Cap Absorption and Related Geomagnetic Effect, J. Geophys. Res. 66, 1594 (1961).

Jelly, D.H. and C. Collins. Some Observations of Polar Cap Absorption in the Northern and Southern Hemispheres, Can. J. Phys. 40, 706 (1962).

Paghis, I. On the Selection of Solar Flare Radio Burst Events that are Associated with Polar Cap Absorption, J. Geophys. Res. 67, 4503 (1962).

Warwick, C.S. and M.W. Haurwitz. A Study of Solar Activity Associated with Polar Cap Absorption, J. Geophys. Res. 67, 1317 (1962).

Kanellakos, D.P., K.L. Chan, and O.G. Villard. On the Altitude at Which Some Solar Flare-Induced Ionization is Released, J. Geophys. Res. 67, 1795 (1962).

Herman, J.R. Polar Cap and Auroral Zone Absorption Effects on 2.5 and 5.0 Megacycles per Second Atmospheric Radio Noise, J. Geophys. Res. 67, 2299 (1962).

Barry, G.H. and P.R. Widess. The Effect of a Solar Flare on the Frequency of High Frequency Ground Backscatter, J. Geophys. Res. 67, 2707 (1962).

Leinbach, H. Interpretations of the Time Variations of Polar Cap Absorption Associated with Solar Cosmic Ray Bombardments. Scientific Report No. 3, University of Alaska (May 1962).



Prince, H. D. and E. R. Hedeman. Studies of Flares Associated with Polar Cap Absorption, Astron. J. 68, 290 (1963).

Warwick, C. S. Delay Time of Polar Cap Blackout and Its Relation to Delay Time of Geomagnetic Disturbance, J. Geophys. Res. 68, 1561 (1963).

Jelly, D. H. The Effects of Polar Cap Absorption on H. F. Oblique - Incidence Circuits, J. Geophys. Res. 68, 1705 (1963).

Khachalana, G. M. Anomalous Absorption in the Polar Cap Resulting from Large Chromospheric Flares, Geomagnetism and Aeronomy 3, 735 (1963).

Bell, B. Type IV Solar Radio Bursts, Geomagnetic Storm, and Polar Cap Absorption (PCA) Events, Smithsonian Inst., Washington (1963).

Hill, G. E. Polar Cap and Auroral Zone Absorption Events During the First 6 Months of the IGY, J. Phys. Soc. Japan, 17, Supplement A-1, International Conference on Cosmic Rays and the Earth Storm, Part 1 (1963).

Masley, A. J. and A. D. Goedeke. Polar Observations During the 1962 and 1963 Solar Cosmic Ray Events, International Conference on Cosmic Rays, 8th, Jaipur, India (2-14 December 1963) Proceedings. Vol. 1, Solar Particles and Sun-Earth Relations.

Bailey, D. K. Polar Cap Absorptions, Planet. Space Sci. 12, 495 (1964).

Weir, R. A. and R. R. Brown. On the Contribution of Solar Flare Alpha Particles to Polar Cap Absorption Events, J. Geophys. Res. 69, 2193 (1964).

Landmark, B. and J. Ortner. Report to Working Group II from the Panel on Polar Cap Experiments, COSPAR Meeting, 7th, and International Space Sciences Symposium, 5th, Florence, Italy (8-20 May 1964).

Van Allen, J. A., W. C. Lin, and H. Leinbach. On the Relationship Between Absolute Solar Cosmic Ray Intensity and Riometer Absorption, J. Geophys. Res. 69, 4481 (1964).

Haurwitz, M. W., S. Yoshida, and S. I. Akasofu. Interplanetary Magnetic Field Asymmetries and Their Effects on Polar Cap Absorption Events and Forbush Decreases, J. Geophys. Res. 70, 2977 (1965).



Leinbach, H., D. Venkatisan, and R. Parthasanathy. The Influence of Geomagnetic Activity on Polar Cap Absorption, State Univ. of Iowa, Iowa City, Dept. of Physics, Astronomy (April 1965).

Michel, F.E. and A.J. Dessler. Physical Significance of Inhomogeneities in Polar Cap Absorption Events, J. Geophys. Res. 70, 4305 (1965).

GEOMAGNETIC DISTURBANCES RELATED TO SOLAR FLARES

Muller, R. et al. Solar and Terrestrial Changes Associated with the Mogel-Dillinger Effects (SID), J. Atmos. Terres. Phys. 1, 37 (1950).

Sarabhai, V. and R.P. Kane. Effects of Geomagnetic Field on Solar Cosmic Rays, Phys. Rev. 92, 415 (1953).

Grahm, J.H. and S.E. Forbush. Solar Flare and Magnetic Storms Effects in Cosmic Ray Intensity Near Geomagnetic North Pole. Phys. Rev. 98, 1348 (1955).

Jory, F.S. Selected Cosmic Ray Orbits in the Earth's Magnetic Field, Phys. Rev. 103, 1068 (1956).

Finch, H.F. and B.R. Leaton. The Earth's Magnetic Field-Epoch 1955.0, Mon. Not., Roy. Astron. Soc., Geophys. Supplement 7, 314 (1957).

Lüst, R. Impact Zones for Solar Cosmic Ray Particles, Phys. Rev. 105, 1827 (1957).


Winckler, J.R. and L. Peterson. Large Auroral Effect on Cosmic Ray Detectors Observed at 8 gm/cm² Atmospheric Depth, Phys. Rev. 108, 903 (1957).

Schlüter, A. Störmer Orbits of Low Energy, Nuovo Cimento, Series X, Supplement 8, 349 (1959).

Lüst, R. Impact Zones for Solar Cosmic Ray Particles, Nuovo Cimento, Series X, Supplement 8, 176 (1958).

Quenby, J.J. and W.R. Webber. Cosmic Ray Cut-Off Rigidities and the Earth's Magnetic Field, Phil. Mag. 4, 90 (1959).

Warwick, C.S. and R.T. Hansen. Geomagnetic Activity Following Large Solar Flares, J. Atmos. Terres. Phys. 14, 287 (1959).



- Bachelet, F. et al. Cosmic Ray and Geomagnetic Disturbances from July 1957 to July 1958 Part I - The Cosmic Ray Events and Their Correlation with Geomagnetic Events, Nuovo Cimento, Series X, 16, 292 (1960).
- Bachelet, F. et al. Cosmic Ray and Geomagnetic Disturbances from July 1957 to July 1958 Part II - The Geomagnetic Events, Nuovo Cimento, Series X, 16, 320 (1960).
- Anderson, K.A. and D.C. Enemark. Observations of Solar Cosmic Rays Near the North Magnetic Pole, J. Geophys. Res. 65, 2657 (1960).
- Kelsall, T. Solar Proton Impact Zones, J. Geophys. Res. 66, 4047 (1961).
- Smith, E. v. P. and P.S. McIntosh. On the Association Between Noise Storms and Solar Flares, J. Geophys. Res. 67, 1013 (1962).
- Keppler, E. et al. Sudden Increase in Radiation Intensity Coinciding with a Geomagnetic Storm Sudden Commencement, J. Geophys. Res. 67, 5343 (1962).
- Pearce, L.S. Frequency and Magnitude of Solar Cosmic-Ray Events as Measured within the Geomagnetic Field, Proceedings of the Lunar and Planetary Exploration Colloq., Santa Monica, Calif., (23-24 May 1962), pp. 63-74.
- Belrose, J.S. and E. Cetiner. Measurement of Electron Densities in the Ionospheric D. Region at the Time of a 2+ Solar Flare, Nature 195, 688 (1962).
- Anthony, R.W., Jr. A Study of Ionospheric Absorption Produced by Solar Cosmic Rays, University of California, Berkeley (1962).
- Kane, R.P. Recovery Times of Geomagnetic Storms, Nuovo Cimento, Series X, 29, 826 (1963).
- Jenkins, R.W. and I. Paghis. Criteria for the Association of Solar Flares with Geomagnetic Disturbances, Can. J. Phys. 41, 1056 (1963).
- Webber, W.R. Effective Cutoff Rigidities for Solar Cosmic Rays in the Earth's Field, Trans. Am. Geophys. Union 44, 72 (1963).
- Sauer, H.H. A New Method of Computing Cosmic Ray Cutoff Rigidity for Several Geomagnetic Field Models, J. Geophys. Res. 68, 957 (1963).
- Akasofu, S.I. and W.C. Lin. The Magnetic Moment of Model Ring Current Belts and the Cutoff Rigidity of Solar Protons, J. Geophys. Res. 68, 973 (1963).



Gregory, J. B. Particle Influx at High Latitudes 3 - Solar Protons, J. Geophys. Res. 68, 3097 (1963).

Lipshitz, M. and A. M. Lenchek. Impulsive Injection of Electrons into the Earth's Inner Trapping Region from Solar Cosmic Ray Events, J. Geophys. Res. 68, 4081 (1963).

Davis, T. N. Negative Correlation Between Polar Cap Visual Aurora and Magnetic Activity, J. Geophys. Res. 68, 4447 (1963).

Akasofu, S. I., W. C. Lin, and J. A. Van Allen. The Anomalous Entry of Low Rigidity Solar Cosmic Rays into the Geomagnetic Field, J. Geophys. Res. 68, 5327 (1963).

Dorman, L. I. and E. V. Kolomeets. The Effect of Small Individual Solar Bursts of Cosmic Rays, (Results of Researches on the Programme of the International Geophysical Year. Section VII), Trans. Acad. Sci. USSR, p. 142 (1963).

Zevakina, R. A. Space-Time Distribution of an Ionospheric Disturbance at High Latitudes of the Northern and Southern Hemispheres After Large Chromospheric Flares, Geomagnetism and Aeronomy 3, 62 (1963).

Nestorav, G., L. Krzhioski, and V. Letfus. Ionospheric Anomalies as an Indicator of Ionizing Radiation of a Corpuscular Cloud After a Solar Chromospheric Flare, Geomagnetizm i Aeronomiia 4, 1059 (1964).

Larmore, L. Recent Observations and Effects of Solar Flares, in International Symposium on Space Technology and Science, Tokyo, 27-31 August 1962, 4th Proceedings, edited by Tamiya Nomuna. Japan Publications Trading Co., Rutland, Vt. (1963).

Ramanathan, K. R. Ionospheric Phenomena and Solar Activity, International Conference on Cosmic Rays, 8th, Jaipur, India, 2-14 December 1963, Proceedings. Vol. 1, Solar Particles and Sun-Earth Relations, pp. 25-38.

Webber, W. R. and F. B. McDonald. Cerenkov Scintillation Counter Measurements of the Intensity and Modulation of Low Rigidity Cosmic Rays and Features of the Geomagnetic Cutoff Rigidity, J. Geophys. Res. 69, 3097 (1964).

Byrne, F. N., M. A. Ellison, and J. H. Reid. A Survey of Solar Flare Phenomena, Space Science Rev. III, 319 (1964).

Volland, H. On the Solar Flare Effect of V. L. F. - Waves in the Lower Ionosphere, J. Atmos. Terres. Phys. 26, 695 (1964).



Brown, R.R. On the Relationship Between Polar Slow Aurora and Solar Cosmic Ray Fluxes, J. Atmos. Terres. Phys. 26, 805 (1964).

Dodson, H.W. and E.R. Hedeman. Problems of Differentiation of Flares with Respect to Geophysical Effects, Planet. Space Sci. 12, 393 (1964).

Gachichiladze, R.S. and G.M. Khochalaya. The State of Ionosphere at Middle Latitudes During Magnetic Storms Associated with an Anomalous Absorption at the Polar Cap, Geomagnetizm i Aeronomiia 4, 602 (1964).

Bateman, R.G. Geomagnetic Effects of Plage Areas on the Sun, Astron. J. 70, 132 (1965).

Zmuda, A.J., G.F. Pieper, and C.O. Bostrom. Trapped Protons in the South Atlantic Magnetic Anomaly, July through December 1961-3, Magnetic Storms and Solar Proton Events, J. Geophys. Res. 70, 2045 (1965).

Catalogue of Disturbances in Ionosphere Geomagnetic Field, Field Intensity and Radio Wave, Cosmic Ray, Solar Phenomena and Other Related Phenomena, Report of Ionosphere and Space Res. in Japan 19, No. 1, pp. 61-119 (March 1965).

Chivers, H.J.A. and J.K. Hargraves. Conjugate Observations of Solar Proton Events Delayed Ionospheric Changes During Twilight, Planet. Space Sci. 13, 583 (1965).

Reid, G.C. Solar Cosmic Rays and the Ionosphere, in Physics of the Earth's Upper Atmosphere, edited by C.O. Hines et al, Prentice Hall, Inc., Englewood Cliffs, N.J. (1965), pp. 245-270.

IONIZATION AND REACTIONS IN THE EARTH'S ATMOSPHERE

Lord, J.J. and M. Schein. A Fluctuation in the Nuclear Disintegration Rate in Photographic Emulsions Exposed in the Stratosphere Following a Solar Flare, Phys. Rev. 78, 321 (1950).

Warwick, C.S. Flare Height and Association with SID's, Astrophys. J. 121, 385 (1955).

Little, C.G. and H. Leinbach. Some Measurements of High-Latitude Ionospheric Absorption Using Extraterrestrial Radio Waves, Proc. IRE 46, 334 (1958).

Bailey, D.K. Abnormal Ionization in the Lower Ionospheric Associated with Cosmic-Ray Flux Enhancements, Proc. IRE 47, 255 (1959).



Little, C.G. and H. Leinbach. The Riometer - A Device for the Continuous Measurement of Ionospheric Absorption, Proc. IRE 47, 315 (1959).

Palmeira, R.A.R. and K.G. McCracken. Observation of a Short-Lived Cosmic Ray Solar Flare Increase with a High Counting Rate Meson Detector, Phys. Rev. Ltrs. 5, 15 (1960).

Simpson, J.A. The Production of Tritons and C14 in the Terrestrial Atmosphere by Solar Protons, J. Geophys. Res. 65, 1615 (1960).

Hill, G.E. Ionospheric Disturbances Following a Solar Flare, J. Geophys. Res. 65, 3183 (1960).

Brown, R.R. and R.A. Weir. Ionospheric Effects of Solar Protons, Arkiv for Geofysik 3, 523 (1961).

Hill, G.E. Effects of Corpuscular Emissions on the Polar Ionosphere Following Solar Flares, J. Geophys. Res. 66, 2329 (1961).

Bhavsar, P.D. Gamma Rays from the Solar Cosmic Ray Produced Nuclear Reactions in the Earth's Atmosphere and Lower Limit on the Energy of Solar Protons Observed at Minneapolis, J. Geophys. Res. 67, 2627 (1962).

Schaefer, H.J. Depth of Penetration of Solar Protons into the Atmosphere and Related Radiation Exposure in Supersonic Transport, Aerospace Medicine 34, 1 (1963).

Chilton, C.J., F.K. Steele, and R.B. Norton. Very Low Frequency Phase Observations of Solar Flare Ionization in the D Region of the Ionosphere, J. Geophys. Res. 68, 5421 (1963).

Webber, W.R. and P.S. Freier. Electron Density Profiles in the Lower Ionosphere Due to Solar Cosmic Rays, J. Geophys. Res. 68, 6223 (1963).

Winkelman, R.E. and R.B. Dyce. Solar Flare Enhancement of Ionization at F Region Heights, J. Geophys. Res. 69, 3625 (1964).

Gauger, J. Anomalous Increase in the Neutron Flux at Aircraft Altitude, J. Geophys. Res. 70, 3571 (1965).

MISCELLANEOUS

Warwick, C.S. Some Characteristics of Solar Flares, Astrophys. J. 120, 237 (1954).

Simpson, J.A. and K.B. Fenton. The Neutron Monitor, University of Chicago Report (1955).



Peterson, L. and J.R. Winckler. Short Gamma-Ray Burst from a Solar Flare, Phys. Rev. Ltrs. 1, 205 (1958).

Van Allen, J.A., C.E. McIlwain, and G.H. Ludwig. Radiation Observations with Satellite 1958 α , J. Geophys. Res. 64, 271 (1959).

Rothwell, P. and C.E. McIlwain. Satellite Observations of Solar Cosmic Rays, Nature 184, 138 (1959).

Peterson, L. and J.R. Winckler. Gamma-Ray Burst from a Solar Flare, J. Geophys. Res. 64, 697 (1959).

Chubb, T.A., H. Friedman, and R.W. Kreplin. Measurements Made of High Energy X-Rays Accompanying Three Class 2 + Solar Flares, J. Geophys. Res. 65, 1831 (1960).

Boischot, A. The Relation Between Radioelectric Emission and Solar Cosmic Rays, in Proceedings of the 11th International Astronautical Congress, Stockholm, 15-20 August 1960, Vol. 1, Main Sessions, I. International Astronautical Federation, Stockholm, Ahlin and Okerbund Civillryckin (1960), pp. 227-231.

Vette, J.I. and F.G. Casal. High Energy X-Rays During Solar Flares, Phys. Rev. Ltrs. 6, 334 (1961).

Maeda, K. and V.L. Patel. A Note on Solar Flare Cosmic Rays, J. Geophys. Res. 66, 1288 (1961).

Lin, W.C. Observation of Galactic and Solar Cosmic Rays from October 13, 1959 to February 17, 1961 with Explorer VII (Satellite 1959 Iota), State University of Iowa Report SUI-61-16 (1961).

Keith, J.E. and A.L. Turkevich. Radioactivity Induced in Discoverer 17 by Solar Flare Protons, J. Geophys. Res. 67, 4525 (1962).

Waddington, C.J. The Hazard of Corpuscular Solar Radiation to Manned Spaceflight, J. Brit. Interplanet. Soc. 18, 277 (1962).

Freier, P.S. and C.J. Waddington. Energetic Deuterons and Tritons Produced by Solar Flares, International Conference on Cosmic Rays, 8th, Jaipur, India, (2-14 December 1963) Proceedings. Vol. 1, Solar Particles and Sun-Earth Relations, pp. 139-141.

Biswas, S. The Composition of Solar Particle Radiation, International Conference on Cosmic Rays, 8th, Jaipur, India (2-14 December 1963), Proceedings. Vol. 1, Solar Particles and Sun-Earth Relations, pp. 43-67.



Earl, J. A. Cloud Chamber Observations of Galactic and Solar Cosmic Ray Electrons, University of Minnesota Report, Minneapolis (1963).

Chasson, R. L. Remarks on Two Recent Papers on Solar Cosmic Ray Events, J. Geophys. Res. 68, 5097 (1963).

Webber, W. R. A Review of Solar Cosmic Ray Events, AAS-NASA Symposium on the Physics of Solar Flares, NASA-SP-50, p. 215 (1963).

Dale, S. H. Radiation Environment for Manned Spacecraft, Institute of Environmental Sciences, 1963 Annual Technical Mtg., Mt. Prospect, Ill. Proceedings, p. 46 (1963).

Simpson, J. A. Recent Investigations of the Low Energy Cosmic and Solar Particle Radiations, Pontifical Academy of Sciences, Scripto Vario, No. 25 (1963).

Webber, W. R. An Evaluation of the Radiation Hazard Due to Solar Particle Events, Report D2-90467, The Boeing Company (Dec. 1963).

Schaefer, H. J. Dosimetric Evaluation of the Alpha Flux in Solar Particle Beams, NASA Order R-75, N65-18049, Naval School of Aviation Medicine, Pensacola, Fla. (11 November 1964).

Anand, S. P. S., L. Oster, and S. Safio. Excitation of Lunar Luminescence by Solar Protons, Nature 202, 1097 (1964).

Acton, L. W. Inverse Compton Effect as a Possible Cause of the X-Ray Radiation of Solar Flares, Nature 204, 64 (1964).

de Jager, C. Solar Ultraviolet and X-Ray Radiation, Research in Geophysics, H. Odishaw, ed., Vol. 1, Sun, Upper Atmosphere and Space. MIT Press, Cambridge, Mass. (1964), pp. 1-42.

Keppler, E. Balloon Instrument Measurement of X-Ray and Solar Protons in the Northern Auroral Zone, Springer-Verlag, Berlin (1964).

Vladimirskii, B. M. and A. K. Pankraton. X-Radiation from Solar Flares and Solar Cosmic Rays, Bull. Acad. Sci. USSR, Physical Series, No. 12, p. 1911 (1964).

Sturrock, P. A. and B. Coppi. A New Model of Solar Flares, Nature 204, 61 (1964).



Modisette, J. L. Space Radiation, in Manned Spacecraft Engineering Design and Operation, Fairchild Publications, Inc., New York (1964), pp. 97-104.

Lin, W. C. and J. A. Van Allen. Observation of Solar Cosmic Rays from October 13, 1959 to February 17, 1961 with Explorer VII, Academic Press, Inc., Englewood Cliffs, N. J. (1964), pp. 195-235.

Krimigis, S. M., J. A. Van Allen, and L. A. Frank. Observations with University of Iowa Equipment on Mariner IV, May-October, 1965, University of Iowa Report 65-36 (25 October 1965).

Simpson, J. A. and J. J. O'Gallagher. Summary of Measurements with the Cosmic Ray Telescope on Mariner IV, NASA Quarterly Review (October 1965).

Modisette, J. L., T. W. Vinson, and A. C. Hardy. Model Solar Proton Environments for Manned Spacecraft Design, NASA, Manned Spacecraft Center, Houston, Texas (April 1965).

Bryant, D. A. et al. Studies of Solar Protons with Explorers XII and XIV, Astrophys. J. 141, 500 (1965).

Van Allen, J. A. et al. Solar X-Ray Observations of Injun I, J. Geophys. Res. 70, 1639 (1965).

Krivsky, L. Disappearance of the Polar Cap on Mars in 1956 After a Solar Flare with Ejection of Particles, Nature 205, 994 (1965).

Kiepenheuer, K. O. About the Flare Problem, in The Solar Spectrum; Proceedings of the Symposium, Univ. of Utrecht, Utrecht, Netherlands, 26-31 August 1963. D. Reidel Publishing Co., Dordrecht, Netherlands (1965), pp. 240-252.



APPENDIX B. FLARE-PLAGE CORRELATION TABLES

Tables B1 through B14 give values of w_{ijk} , the probability of occurrence of at least one flare (with corrected area 100 millionths of the solar hemisphere or greater) at a calcium plage region with given area, age, and luminosity. Odd-numbered tables refer to flares on the same disk passage as that on which the plage is observed; even-numbered tables refer to flares on the next disk passage of the position of the plage. A pair of tables is presented for each year from 1958 through 1964; these years correspond to 1969 through 1975, in sunspot number cycle 20.

When $P_{ijk} = 0$ in Equation 10, w_{ijk} is indeterminate; i. e., no plages with the given characteristics were observed. These values of w_{ijk} may be interpolated or extrapolated from adjacent values or adjacent years.



Table B1. Probability of Flare Occurrence on Same Rotation,
1958 (1969)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
<1000	1	0.148	0.404	0.800	1
	2	0.136	0.478	0.500	*
	3	0.133	0.556	*	*
	4	0.300	0.500	1	*
	5	0.125	0.667	*	*
	6	0.200	0.667	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
1000-2000	1	0.500	0.533	0.706	1
	2	0.556	0.563	0.714	1
	3	0.125	0.692	0.667	*
	4	0.333	0.667	1	*
	5	0.333	0.625	*	*
	6	0.500	0.833	1	*
	7	*	*	*	*
	≥ 8	*	*	*	*
2000-4000	1	*	0.800	0.944	*
	2	*	0.875	0.909	*
	3	*	0.800	0.875	*
	4	1	0.750	1	*
	5	*	0.800	1	*
	6	*	0.500	1	*
	7	*	1	1	*
	≥ 8	*	*	*	*
≥ 4000	1	*	1	0.971	1
	2	*	1	1	1
	3	*	1	1	*
	4	*	1	1	1
	5	*	1	1	*
	6	*	*	1	*
	7	*	*	*	*
	≥ 8	*	*	*	*

* = indeterminate

Table B2. Probability of Flare Occurrence on Next Rotation,
1958 (1969)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
<1000	1	0.041	0.128	0.522	0
	2	0.100	0.111	0	*
	3	0.071	0	*	*
	4	0.154	0.375	0	*
	5	0.083	0.250	*	*
	6	0	0	*	*
	7	*	*	*	*
	≥ 8	0	*	*	*
1000-2000	1	0.250	0.162	0.120	1
	2	0.200	0.043	0.111	1
	3	0	0.428	0.286	*
	4	0	0.273	0.667	*
	5	0	0.357	0	*
	6	0.333	0.222	1	*
	7	0	0	1	*
	≥ 8	*	*	*	*
2000-4000	1	*	0.412	0.333	*
	2	*	0.304	0.236	*
	3	0	0.236	0.267	*
	4	0	0	0.250	*
	5	1	0	0.500	*
	6	0	0	0	1
	7	*	0	0	*
	≥ 8	0	*	*	*
≥ 4000	1	*	0.200	0.293	0.667
	2	0.500	0.750	0.500	0.500
	3	*	0.444	0.640	*
	4	0	1	0.400	1
	5	*	0	0.400	0
	6	*	1	0	*
	7	*	*	*	*
	≥ 8	*	*	*	*

* = indeterminate



Table B3. Probability of Flare Occurrence on Same Rotation, 1959 (1970)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
<1000	1	0.222	0.431	0.818	*
	2	0.125	0.545	0.500	*
	3	0.250	0.400	*	*
	4	*	1	1	*
	5	0.200	0.500	*	*
	6	*	0.750	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
1000-2000	1	0.500	0.538	0.833	*
	2	*	0.455	0.667	*
	3	1	0.444	1	*
	4	*	0.571	*	*
	5	1	0.600	1	*
	6	1	0.333	*	*
	7	0.500	1	*	*
	≥ 8	*	*	1	*
2000-4000	1	*	0.571	0.952	*
	2	*	0.706	0.700	*
	3	*	0.667	0.700	*
	4	*	0.750	1	*
	5	*	0.750	1	*
	6	*	1	0.750	*
	7	1	0.500	1	*
	≥ 8	*	*	1	*
≥ 4000	1	*	*	1	*
	2	*	1	1	*
	3	*	0.750	0.923	*
	4	*	1	1	1
	5	*	0.500	1	*
	6	*	1	1	*
	7	*	*	*	*
	≥ 8	*	1	*	1

* = indeterminate

Table B4. Probability of Flare Occurrence on Next Rotation,
1959 (1970)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
< 1000	1	0.172	0.159	0.333	*
	2	0.625	*	*	*
	3	0.833	0.267	*	*
	4	*	0.500	1	*
	5	0.143	0.250	*	*
	6	*	0.250	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
1000-2000	1	*	0.154	0.368	*
	2	*	0.280	0.666	*
	3	*	0.250	0.500	*
	4	1	0.400	*	*
	5	*	0.091	*	*
	6	*	0.666	*	*
	7	*	1	*	*
	≥ 8	*	1	*	*
2000-4000	1	*	0.125	0.524	*
	2	*	0.500	0.333	*
	3	*	0.400	0.429	*
	4	*	0.714	0.800	*
	5	*	0.571	1	*
	6	*	*	0.250	*
	7	1	0.333	*	*
	≥ 8	*	*	0.666	*
≥ 4000	1	*	*	0.625	*
	2	*	0.250	0.750	*
	3	*	0.750	0.785	*
	4	*	1	0.889	*
	5	*	0.500	0.750	*
	6	*	0.500	0.666	*
	7	*	0.500	1	*
	≥ 8	*	*	*	*

* = indeterminate



Table B5. Probability of Flare Occurrence on Same Rotation,
1960 (1971)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
<1000	1	0.235	0.286	0.750	*
	2	*	0.235	*	*
	3	0.250	0.250	*	*
	4	0.143	0.250	*	*
	5	0.200	0.111	*	*
	6	*	*	*	*
	7	*	*	1	*
	≥ 8	*	*	*	*
1000-2000	1	0.500	0.529	0.900	*
	2	0.333	0.308	0.333	*
	3	*	0.333	1	*
	4	0.167	0.214	*	*
	5	*	0.727	1	*
	6	*	0.500	*	*
	7	*	*	*	*
	≥ 8	*	0.333	*	*
2000-4000	1	*	1	0.733	*
	2	*	0.846	0.688	*
	3	*	0.667	0.933	*
	4	*	0.556	0.833	*
	5	*	0.800	1	*
	6	*	1	1	*
	7	*	0.667	1	*
	≥ 8	*	0.500	*	*
≥ 4000	1	*	*	1	*
	2	*	1	1	*
	3	*	1	1	*
	4	*	0.500	0.750	*
	5	*	*	1	*
	6	*	*	0.667	*
	7	*	*	1	*
	≥ 8	*	*	1	*

* = indeterminate



Table B6. Probability of Flare Occurrence on Next Rotation,
1961 (1972)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
<1000	1	0.100	0.053	0.411	*
	2	0	0.154	*	*
	3	0	0.222	0.667	*
	4	0	0	*	*
	5	0.667	0.250	*	*
	6	0	0	*	*
	7	*	*	*	*
	≥8	0	0	*	*
1000-2000	1	*	0.167	0.200	*
	2	*	0.222	0.111	*
	3	0	0.357	0.750	*
	4	0.500	0.167	0.333	*
	5	1	0	*	*
	6	0	0.333	*	*
	7	*	0	0	*
	≥8	*	0	*	*
2000-4000	1	*	0	0.400	*
	2	*	0.500	0.143	*
	3	*	0.667	0.500	*
	4	*	*	0.250	*
	5	*	0.500	0	*
	6	*	*	0	*
	7	*	*	1	*
	≥8	*	0	*	*
≥4000	1	*	*	1	*
	2	*	0.500	0.750	*
	3	*	*	0.667	*
	4	*	*	0.500	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	0	*
	≥8	*	*	*	*

* = indeterminate



Table B7. Probability of Flare Occurrence on Same Rotation,
1961 (1972)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
<1000	1	0.111	*	0.529	*
	2	0.200	*	*	*
	3	*	*	0.666	*
	4	0.333	0.333	*	*
	5	*	*	*	*
	6	0.250	0.500	*	*
	7	*	*	*	*
	≥8	*	0.500	*	*
1000-2000	1	*	0.833	0.720	*
	2	*	0.400	0.250	*
	3	*	0.133	0.500	*
	4	*	0.333	1	*
	5	1	0.666	*	*
	6	*	0.666	*	*
	7	*	*	1	*
	≥8	*	*	*	*
2000-4000	1	*	1	0.789	*
	2	*	0.400	0.833	*
	3	*	0.333	0.750	*
	4	*	*	1	*
	5	*	*	0.500	*
	6	*	*	1	*
	7	*	*	*	*
	≥8	*	*	*	*
≥4000	1	*	*	1	*
	2	*	1	1	*
	3	*	*	1	*
	4	*	*	0.500	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥8	*	*	*	*

* = indeterminate

Table B8. Probability of Flare Occurrence on Next Rotation,
1961 (1972)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
< 1000	1	0.100	0.056	0.412	*
	2	*	0.154	*	*
	3	*	0.182	0.666	*
	4	*	*	*	*
	5	0.666	0.250	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥8	*	*	*	*
1000-2000	1	*	0.167	0.200	*
	2	*	0.182	0.111	*
	3	*	0.357	0.750	*
	4	0.500	0.167	0.111	*
	5	1	*	*	*
	6	*	0.333	*	*
	7	*	*	*	*
	≥8	*	*	*	*
2000-4000	1	*	*	0.400	*
	2	*	0.500	0.143	*
	3	*	0.666	0.500	*
	4	*	*	0.250	*
	5	*	0.500	*	*
	6	*	*	*	*
	7	*	*	1	*
	≥8	*	*	*	*
≥4000	1	*	*	1	*
	2	*	0.500	0.750	*
	3	*	*	0.666	*
	4	*	*	0.500	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥8	*	*	*	*

* = indeterminate



Table B9. Probability of Flare Occurrence on Same Rotation,
1962 (1973)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
< 1000	1	*	*	*	*
	2	*	*	*	*
	3	*	*	*	*
	4	*	*	*	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
1000-2000	1	*	0.300	0.650	*
	2	*	0.333	0.636	*
	3	*	*	0.600	*
	4	*	0.167	*	*
	5	*	0.500	*	*
	6	*	0.333	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
2000-4000	1	0.105	0.036	0.444	*
	2	*	*	0.900	*
	3	1	0.333	1	*
	4	*	*	0.333	*
	5	*	*	0.666	*
	6	*	*	1	*
	7	*	*	*	*
	≥ 8	*	*	*	*
≥ 4000	1	0.250	0.239	0.400	1
	2	0.111	0.300	0.400	*
	3	*	0.111	1	*
	4	*	*	1	*
	5	*	0.500	1	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*

* = indeterminate

Table B10. Probability of Flare Occurrence on Next Rotation,
1962 (1973)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
< 1000	1	*	*	*	*
	2	*	*	*	*
	3	*	*	*	*
	4	*	*	*	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
1000-2000	1	*	*	0.087	*
	2	*	*	*	*
	3	*	*	*	*
	4	*	*	1	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
2000-4000	1	*	*	0.111	*
	2	*	*	*	*
	3	*	*	0.125	*
	4	*	*	0.250	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
≥ 4000	1	*	*	*	*
	2	*	*	*	*
	3	*	*	*	*
	4	*	*	*	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	0.500	1	*	*

* = indeterminate

Table B11. Probability of Flare Occurrence on Same Rotation,
1963 (1974)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
< 1000	1	0.017	0.039	0.200	*
	2	*	*	*	*
	3	*	*	*	*
	4	*	0.143	*	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
1000-2000	1	*	0.250	0.929	1
	2	*	0.166	0.400	*
	3	*	*	0.400	*
	4	*	*	*	*
	5	*	*	*	*
	6	*	*	1	*
	7	*	*	*	*
	≥ 8	*	*	*	*
2000-4000	1	*	0.043	0.857	*
	2	*	*	0.714	*
	3	*	*	0.800	*
	4	*	*	0.666	*
	5	*	1	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
≥ 4000	1	0.083	0.100	0.500	*
	2	*	0.250	1	1
	3	*	*	*	*
	4	*	*	1	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*

* = indeterminate



Table B12. Probability of Flare Occurrence on Next Rotation,
1963 (1974)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
<1000	1	0.013	0.055	*	*
	2	*	*	*	*
	3	*	*	*	*
	4	*	0.143	*	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
1000-2000	1	*	0.125	0.176	1
	2	*	*	0.167	*
	3	*	0.200	*	*
	4	*	*	*	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
2000-4000	1	*	*	*	*
	2	*	0.500	0.222	*
	3	*	*	0.600	*
	4	*	*	*	*
	5	*	1	0.500	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
≥ 4000	1	*	0.048	0.111	*
	2	*	*	0.500	1
	3	*	*	*	*
	4	*	*	1	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*

* = indeterminate



Table B13. Probability of Flare Occurrence on Same Rotation,
1964 (1975)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
< 1000	1	0.016	0.069	0.419	*
	2	0.111	0.333	*	*
	3	*	*	*	*
	4	*	*	*	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
1000-2000	1	*	0.500	0.714	*
	2	*	0.333	0.500	*
	3	*	0.666	1	*
	4	*	*	1	*
	5	*	*	1	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
2000-4000	1	*	1	1	*
	2	*	*	1	*
	3	*	*	*	*
	4	*	*	*	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
≥ 4000	1	*	*	*	*
	2	*	*	*	*
	3	*	*	*	*
	4	*	*	*	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*

* = indeterminate



Table B14. Probability of Flare Occurrence on Next Rotation,
1964 (1975)

Plage Area	Plage Age	Plage Luminosity			
		1-1.5	2-2.5	3-3.5	4-5
< 1000	1	0.003	0.028	0.094	*
	2	*	*	*	*
	3	*	*	*	*
	4	*	*	*	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
1000-2000	1	*	*	0.250	*
	2	*	*	0.333	*
	3	*	0.333	*	*
	4	*	*	1	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
2000-4000	1	*	*	*	*
	2	*	*	0.500	*
	3	*	*	*	*
	4	*	*	1	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*
≥ 4000	1	*	*	*	*
	2	*	*	*	*
	3	*	*	*	*
	4	*	*	*	*
	5	*	*	*	*
	6	*	*	*	*
	7	*	*	*	*
	≥ 8	*	*	*	*

* = indeterminate



APPENDIX C. COMPUTER PROGRAMS

One computer program listed and described below was prepared in this study. Other programs used in the study were completed, under the sole support of North American Aviation, Inc., prior to the inception of this contract.

DESCRIPTION OF PROGRAM

Name: SPOTNO

Purpose: This program calculates the monthly smoothed Wolf sunspot number R_M at each month from a given date to a date 250 years later.

Method: The program evaluates the sum of the contributions to R_M of each of 1 to 50 terms which are a discrete representation of a power frequency spectrum $P(f)$ of the R_M from July 1749 to June 1964. $P(f)$ is defined by Equation 3 in this report, and the summation process replaces the integral in Equation 4 by

$$R_M(t'') = \sum_{n=1}^N P(f_n) (f_{n+1} - f_n) \sin 2\pi f_n (t'' - t_0) \quad (C-1)$$

where t_0 is a time at which $R_M(t_0) = 0$.

Language: FORTRAN 4, for the IBM 7094 computer; compiler - FAP; operating system - IBSYS.

Instructions for Use: Compilation time - 7 seconds; loading time - 17 seconds; execution time - 10 seconds; printing - 3000 lines. Program requires subroutine DECRD, which is on NAA library tapes.

Program Decks: Source, BCD, 29 cards; object - relocatable column binary, 16 cards. These decks are transmitted separately. Program is on file at NAA as Deck 3M906.

Input-Output Variables: FORTRAN variables appearing in input and output statements of SPOTNO are defined in Table C1, in the order of their appearance in the program.




Table C1. Definition of Variables in SPOTNO

FORTRAN Name	Algebraic Symbol	Definition
TO	t_0	Time at which sunspot number = 0 (year AD)
T1	$t''(\text{min})$	Earliest date at which sunspot number is to be found (year AD)
NF	N	Number of terms in power frequency spectrum, Equation C1
F(I)	f_n	Frequency of nth term (year^{-1})
P(I)	$P(f_n)$	Amplitude of nth term
TC	t''	Any date at which sunspot number is found (year AD)
SN	R_M	Sunspot number

LISTING

The listing of SPOTNO follows (Figure C1).

DATA

Values of data variables used in this study are given in Table C2.



```

SPOTNO
SPOTPS      - EFN      SOURCE STATEMENT - IFN(S) -
03/11/86

C      COMPUTES EXPECTED SUNSPOT WOLF NUMBER AT TIME TP FROM POWER
C      SPECTRUM OF FREQUENCIES BASED ON ANNUAL SMOOTHED WOLF NUMBER
C      (1749-1961). F = CYCLES PER YEAR, FO = MAXIMUM F, P = POWER AT F,
C      TP = TIME (YR) AFTER 1 JULY 1749 AT WHICH SN = SUNSPOT WOLF
C      NUMBER IS TO BE FOUND. TC = CALENDAR DATE OF TP, TO = DATE AT
C      WHICH SN = 0. (1810), NF = NUMBER OF VALUES OF F, P.
C      THIS VERSION DOES EACH MONTH JUL 1749 - JUN 1999. T1 = 1ST DATE.
C      DIMENSION F(50), P(50), DA(2)
C      EQUIVALENCE (DA(1), TO), (DA(2), T1)
10 READ (5,710) NF, (F(I), P(I), I = 1, NF)
710 FORMAT (1I12/(2E12.5))
20 CALL DECRO(DA)
   TO = TO - 1749.0
   FO = F(NF)
810 WRITE (6,910)
910 FORMAT (17H1 DATE SPOTS)
22 DO 200 IT = 1, 3000
24 TC = T1 + FLGAT(IT - 1)/12.0
30 TP = TC - 1749.0
100 XSUM = 0.0
   NF1 = NF - 1
110 DO 150 I = 1, NF1
120 XSUM = XSUM + P(I) * (F(I+1) - F(I)) * SIN (6.2832 * F(I) * (TP
   1 - TO))
150 CONTINUE
160 SN = SQRT(ABS(XSUM)/FO)
820 WRITE (6,920) TC, SN
920 FORMAT (1H0,F8.2,2X,F6.1)
200 CONTINUE
   GO TO 20
END
SSPS0015
SSPS0020
SSPS0025
SSPS0030
SSPS0035
SSPS0040
SSPS0041
SSPS0045
SSPS0047
SSPS0050
SSPS0055
SSPS0058
SSPS0060
SSPS0062
SSPS0065
SSPS0070
SSPS0076
SSPS0078
SSPS0080
SSPS0085
SSPS0090
SSPS0095
SSPS0100
SSPS0101
SSPS0105
SSPS0110
SSPS0115
SSPS0120
SSPS0122
SSPS0125
SSPS0130

```

1

11

14

29

34

35

Figure C1. SPOTNO Listing



Table C2. SPOTNO Data.

$t_o = 1810.5$ $t'' \text{ (min)} = 1749.5$ $n = 25$		
n	$f_n (\text{year}^{-1})$	$P(f_n) (\text{spots}^2/\text{yr}^2)$
1	0.0045	4,600
2	0.0112	21,500
3	0.0176	12,300
4	0.0233	6,100
5	0.0281	700
6	0.0345	2,200
7	0.0406	1,200
8	0.0475	3,400
9	0.0522	600
10	0.0567	2,100
11	0.0650	2,200
12	0.0705	4,800
13	0.0765	5,200
14	0.0828	5,200
15	0.0903	41,700
16	0.0945	10,500
17	0.1011	30,200
18	0.1063	15,300
19	0.1122	5,600
20	0.1178	7,800
21	0.1236	5,200
22	0.1290	1,300
23	0.1340	2,100
24	0.1410	1,500
25	0.1450	0